



Cochrane
Library

Cochrane Database of Systematic Reviews

18F PET with florbetaben for the early diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI) (Review)

Martínez G, Vernooij RWM, Fuentes Padilla P, Zamora J, Flicker L, Bonfill Cosp X

Martínez G, Vernooij RWM, Fuentes Padilla P, Zamora J, Flicker L, Bonfill Cosp X.
18F PET with florbetaben for the early diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI).
Cochrane Database of Systematic Reviews 2017, Issue 11. Art. No.: CD012883.
DOI: [10.1002/14651858.CD012883](https://doi.org/10.1002/14651858.CD012883).

www.cochranelibrary.com

18F PET with florbetaben for the early diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI) (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	7
OBJECTIVES	10
METHODS	10
RESULTS	12
Figure 1.	13
Figure 2.	15
Figure 3.	16
DISCUSSION	17
AUTHORS' CONCLUSIONS	19
ACKNOWLEDGEMENTS	19
REFERENCES	21
CHARACTERISTICS OF STUDIES	26
DATA	33
Test 1. 18F-florbetaben visual assessment and progression to ADD.	33
Test 2. 18F-florbetaben SUVR and progression to ADD.	33
Test 3. 18F-florbetaben visual assessment and progression to any other form of non-ADD.	33
Test 4. 18F-florbetaben SUVR and progression to any other form of non-ADD.	33
Test 5. 18F-florbetaben visual assessment and progression to any form of dementia.	33
Test 6. 18F-florbetaben SUVR and progression to any form of dementia.	33
APPENDICES	33
CONTRIBUTIONS OF AUTHORS	42
DECLARATIONS OF INTEREST	43
SOURCES OF SUPPORT	43
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	43

[Diagnostic Test Accuracy Review]

18F PET with florbetaben for the early diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI)

Gabriel Martínez^{1,2,3}, Robin WM Vernooij¹, Paulina Fuentes Padilla^{1,2}, Javier Zamora⁴, Leon Flicker⁵, Xavier Bonfill Cosp^{6,7}

¹Iberoamerican Cochrane Centre, Barcelona, Spain. ²Faculty of Medicine and Dentistry, Universidad de Antofagasta, Antofagasta, Chile.

³Alzheimer Research Center and Memory Clinic of Fundació ACE, Institut Català de Neurociències Aplicades, Barcelona, Spain. ⁴Clinical Biostatistics Unit, Ramon y Cajal Institute for Health Research (IRYCIS), CIBER Epidemiology and Public Health (CIBERESP), Madrid (Spain) and Women's Health Research Unit, Centre for Primary Care and Public Health, Queen Mary University of London, London, UK.

⁵Western Australian Centre for Health & Ageing - WACHA, University of Western Australia, Perth, Australia. ⁶Iberoamerican Cochrane Centre, Biomedical Research Institute Sant Pau (IIB Sant Pau), CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain.

⁷Universitat Autònoma de Barcelona, Barcelona, Spain

Contact address: Gabriel Martínez, Iberoamerican Cochrane Centre, C/ Sant Antoni Maria Claret 167, Pavelló 18 Planta 0, Barcelona, Barcelona, 08025, Spain. gmartinez@cochrane.es, gmartinezfuentes@gmail.com.

Editorial group: Cochrane Dementia and Cognitive Improvement Group.

Publication status and date: New, published in Issue 11, 2017.

Citation: Martínez G, Vernooij RWM, Fuentes Padilla P, Zamora J, Flicker L, Bonfill Cosp X. 18F PET with florbetaben for the early diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI). *Cochrane Database of Systematic Reviews* 2017, Issue 11. Art. No.: CD012883. DOI: [10.1002/14651858.CD012883](https://doi.org/10.1002/14651858.CD012883).

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

¹⁸F-florbetaben uptake by brain tissue, measured by positron emission tomography (PET), is accepted by regulatory agencies like the Food and Drug Administration (FDA) and the European Medicine Agencies (EMA) for assessing amyloid load in people with dementia. Its added value is mainly demonstrated by excluding Alzheimer's pathology in an established dementia diagnosis. However, the National Institute on Aging and Alzheimer's Association (NIA-AA) revised the diagnostic criteria for Alzheimer's disease and confidence in the diagnosis of mild cognitive impairment (MCI) due to Alzheimer's disease may be increased when using some amyloid biomarkers tests like ¹⁸F-florbetaben. These tests, added to the MCI core clinical criteria, might increase the diagnostic test accuracy (DTA) of a testing strategy. However, the DTA of ¹⁸F-florbetaben to predict the progression from MCI to Alzheimer's disease dementia (ADD) or other dementias has not yet been systematically evaluated.

Objectives

To determine the DTA of the ¹⁸F-florbetaben PET scan for detecting people with MCI at time of performing the test who will clinically progress to ADD, other forms of dementia (non-ADD), or any form of dementia at follow-up.

Search methods

The most recent search for this review was performed in May 2017. We searched MEDLINE (OvidSP), Embase (OvidSP), PsycINFO (OvidSP), BIOSIS Citation Index (Thomson Reuters Web of Science), Web of Science Core Collection, including the Science Citation Index (Thomson Reuters Web of Science) and the Conference Proceedings Citation Index (Thomson Reuters Web of Science), LILACS (BIREME), CINAHL (EBSCOhost), ClinicalTrials.gov (<https://clinicaltrials.gov>), and the World Health Organization International Clinical Trials Registry Platform (WHO ICTRP) (<http://www.who.int/ictpr/search/en/>). We also searched ALOIS, the Cochrane Dementia & Cognitive Improvement Group's specialised register of dementia studies (<http://www.medicine.ox.ac.uk/alosis/>). We checked the reference lists of any relevant studies

and systematic reviews, and performed citation tracking using the Science Citation Index to identify any additional relevant studies. No language or date restrictions were applied to electronic searches.

Selection criteria

We included studies that had prospectively defined cohorts with any accepted definition of MCI at time of performing the test and the use of ^{18}F -florbetaben scan to evaluate the DTA of the progression from MCI to ADD or other forms of dementia. In addition, we only selected studies that applied a reference standard for Alzheimer's dementia diagnosis, for example, the National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association (NINCDS-ADRDA) or Diagnostic and Statistical Manual of Mental Disorders-IV (DSM-IV) criteria.

Data collection and analysis

We screened all titles and abstracts identified in electronic-database searches. Two review authors independently selected studies for inclusion and extracted data to create two-by-two tables, showing the binary test results cross-classified with the binary reference standard. We used these data to calculate sensitivities, specificities, and their 95% confidence intervals. Two independent assessors performed quality assessment using the QUADAS-2 tool plus some additional items to assess the methodological quality of the included studies.

Main results

Progression from MCI to ADD, any other form of dementia, and any form of dementia was evaluated in one study (Ong 2015). It reported data on 45 participants at four years of follow-up; 21 participants met NINCDS-ADRDA criteria for Alzheimer's disease dementia at four years of follow-up, the proportion converting to ADD was 47% of the 45 participants, and 11% of the 45 participants met criteria for other types of dementias (three cases of Frontotemporal Dementia (FTD), one of Dementia with Lewy body (DLB), and one of Progressive Supranuclear Palsy (PSP)). We considered the study to be at high risk of bias in the domains of the reference standard, flow, and timing (QUADAS-2).

MCI to ADD; ^{18}F -florbetaben PET scan analysed visually: the sensitivity was 100% (95% confidence interval (CI) 84% to 100%) and the specificity was 83% (95% CI 63% to 98%) (n = 45, 1 study). Analysed quantitatively: the sensitivity was 100% (95% CI 84% to 100%) and the specificity was 88% (95% CI 68% to 97%) for the diagnosis of ADD at follow-up (n = 45, 1 study).

MCI to any other form of dementia (non-ADD); ^{18}F -florbetaben PET scan analysed visually: the sensitivity was 0% (95% CI 0% to 52%) and the specificity was 38% (95% CI 23% to 54%) (n = 45, 1 study). Analysed quantitatively: the sensitivity was 0% (95% CI 0% to 52%) and the specificity was 40% (95% CI 25% to 57%) for the diagnosis of any other form of dementia at follow-up (n = 45, 1 study).

MCI to any form of dementia; ^{18}F -florbetaben PET scan analysed visually: the sensitivity was 81% (95% CI 61% to 93%) and the specificity was 79% (95% CI 54% to 94%) (n = 45, 1 study). Analysed quantitatively: the sensitivity was 81% (95% CI 61% to 93%) and the specificity was 84% (95% CI 60% to 97%) for the diagnosis of any form of dementia at follow-up (n = 45, 1 study).

Authors' conclusions

Although we were able to calculate one estimation of DTA in, especially, the prediction of progression from MCI to ADD at four years follow-up, the small number of participants implies imprecision of sensitivity and specificity estimates. We cannot make any recommendation regarding the routine use of ^{18}F -florbetaben in clinical practice based on one single study with 45 participants. ^{18}F -florbetaben has high financial costs, therefore, clearly demonstrating its DTA and standardising the process of the ^{18}F -florbetaben modality are important prior to its wider use.

PLAIN LANGUAGE SUMMARY

^{18}F PET with florbetaben for the early diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment

Review question: In people with mild cognitive impairment (MCI), does using a ^{18}F PET scan with florbetaben predict progression to Alzheimer's disease dementia (ADD) and other dementias?

Background

Due to global ageing, the number of people with dementia is expected to increase dramatically in the next few decades. Diagnosing dementia at an early stage is desirable, but there is no widespread agreement on the best approach. A range of simple pen and paper tests used by healthcare professionals can assess people with poor memory or cognitive impairment. Whether or not using special PET scans that detect amyloid—one of the hallmarks of Alzheimer's disease—improves our ability to predict the progression from MCI to ADD or other forms of dementia remains unclear. Since these tests are expensive, it is important that they provide additional benefits.

Aim

We aimed to evaluate the accuracy of the ^{18}F -florbetaben PET scan in identifying those people with MCI who clinically progress to ADD, other types of dementia, or any form of dementia over a period of time.

Study characteristics

The evidence is current to May 2017. We found 1 study including 45 participants with MCI with a follow-up of 4 years; gender was not reported and the median age for those with a PET-positive scan by quantitative assessment was 73.5 years old. For those with a PET-negative scan the mean age was 71.8 years old. Participants were mainly recruited from local memory clinics.

Study funding sources: the study was funded by the test manufacturer.

Quality of the evidence

The main limitation of this review was that our findings were based on only one study, with not enough details on how the participants were selected. The study was considered to be at high risk of bias, since the final ADD diagnosis was not established separately from the scan results, and due to potential conflicts of interest detected.

Key findings

In this review, based on only one study, we found that the ^{18}F -florbetaben PET scan, as a single test with visual assessment, correctly classified 100% of the participants who will progress to ADD and 83% of the participants who did not progress to ADD at four years follow-up. This means that in a cohort with 100 participants with MCI, 47 of whom will progress to ADD, we would expect that all those 47 MCI participants would test positive with the ^{18}F -florbetaben scan and that 0 participants would be falsely negative (i.e. none of the 47 participants would have a negative test and yet progress to ADD). In addition, we would expect 44 of 53 participants who did not progress to ADD to be ^{18}F -florbetaben-negative and 9 to be falsely positive (i.e. 9 of the 53 participants would have a positive test but not progress to ADD).

The small size of the included study lowered our confidence on these estimates of accuracy and it is still possible that the test is considerably less accurate than these results suggest.

We conclude that ^{18}F -florbetaben imaging is a promising test to predict the progression from MCI to ADD; however, we need more studies to clearly demonstrate its accuracy.

SUMMARY OF FINDINGS

Summary of findings 1. Diagnostic test accuracy of ¹⁸F-florbetaben to predict the progression to ADD, any other form of dementia (non-ADD) or any form of dementia in people with MCI

What is the diagnostic accuracy of ¹⁸F-florbetaben PET amyloid biomarker for predict progression to ADD or any other form of dementia (non-ADD) or any form of dementia in people with MCI?

Descriptive

Patient population	Participants diagnosed with MCI at baseline using any of the Petersen criteria or Winblad criteria or CDR = 0.5 or any 16 definitions included by Matthews (Matthews 2008)
Sources of referral	Memory clinic
MCI criteria	Petersen criteria 2004 and Winblad 2004 (Petersen 2004 ; Winblad 2004)
Sampling procedure	unclear
Prior testing	The only testing prior performing the ¹⁸ F-florbetaben PET amyloid biomarker was the application of diagnostic criteria for identifying participants with MCI
Settings	Secondary care
Index test	¹⁸ F-florbetaben PET
Threshold prespecified at baseline	Yes
Threshold interpretation	Visual and quantitative
Threshold	Visual: if any tracer uptake was visible in any of the frontal, parietal, temporal, and posterior cingulate/precuneus cortices SUVR (Standardised Uptake Volume ratio) of ROI: ≥ 1.45
¹⁸ F-florbetaben retention region	Visual: frontal, parietal, temporal, and posterior cingulate/precuneus cortices Global cortex (SUVR) SUVR: Global cortex
Reference Standard	For Alzheimer's disease dementia: NINCDS-ADRDA (McKhann 1984) For Lewy body dementia:

McKeith criteria ([McKeith 2005](#))

For frontotemporal dementia:

Lund criteria ([Brun 1994](#))

For progressive supranuclear palsy:

Preliminary NINDS criteria ([Hauw 1994](#))

Target condition	Progression from MCI to Alzheimer’s disease dementia or any other forms of dementia or any form of dementia.						
Included studies	Prospectively well-defined cohorts with any accepted definition of MCI (as above). One study (N = 45 participants) was included. Number of participants included in analysis: 45.						
Quality concerns	Patient characteristics were poorly reported. Reference standard diagnosis was made with knowledge of the index test. Applicability concerns were high in reference standard.						
Limitations	We were not able to calculate a summary of sensitivity and specificity due to insufficient number of studies. Investigation of heterogeneity and sensitivity analysis were not done due to insufficient number of studies.						
Test	Studies	Cases/Partic- ipants	Sensitivity	Specificity	Consequences in a cohort of 100		
					Proportion converting ¹	Missed cas- es ²	Overdiag- nosed
Alzheimer's disease dementia							
18F-florbetaben (visual assess- ment)	1	21/45	100% (95% CI 84% to 100%)	83% (95% CI 63% to 95%)	47	0	9
18F-florbetaben (SUVR)	1	21/45	100% (95% CI 84% to 100%)	88% (95% CI 68% to 97%)	47	0	6
Any other form of dementia (non-ADD)							
18F-florbetaben (visual assess- ment)	1	5/45	0% (95% CI 0% to 52%)	38% (95% CI 23% to 54%)	11	11	55
18F-florbetaben (SUVR)	1	5/45	0% (95% CI 0% to 52%)	40% (95% CI 25% to 57%)	11	11	53
Any form of dementia							

¹⁸ F-florbetaben (visual assessment)	1	26/45	81% (95% CI 61% to 93%)	79% (95% CI 54% to 94%)	58	11	9
¹⁸ F-florbetaben (SUVR)	1	26/45	81% (95% CI 61% to 93%)	84% (95% CI 60% to 97%)	58	11	7

Investigation of heterogeneity and sensitivity analysis: The planned investigations of heterogeneity or sensitivity analyses were not possible due to a limited number of studies available for each analysis.

Conclusions: ¹⁸F-florbetaben PET scan has a good sensitivity achieved especially in predicting the progression from MCI to ADD. The quality of evidence was weak because it was based on only one study (45 participants) and there was high risk of bias due to the knowledge of the reference standard to do the diagnosis at four-year follow-up and due to possible conflict of interest detected. There is a need for conducting studies using standardised ¹⁸F-florbetaben PET scan methodology in larger populations. Regarding the aforementioned we do not recommend the use in clinical practice until the DTA performance will be clearly demonstrated.

1. Proportion converting to ADD or any other form of dementia (non-ADD) or any form of dementia in the included study
2. Missed and overdiagnosed numbers were computed using the proportion converting to the target condition.

ADD: Alzheimer's disease dementia

CDR: Clinical Dementia Rating

MCI: Mild cognitive impairment

NINCDS-ADRDA: National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association

NINDS: National Institute of Neurological Disorders and Stroke

PET: Positron emission tomography

ROI: Region of interest

SUVR: Standardised uptake value ratio

BACKGROUND

Dementia is a syndrome due to a brain disease — usually of a chronic or progressive nature — in which there is disturbance of multiple higher cortical functions, including memory, thinking, orientation, comprehension, calculation, learning capacity, language, and judgement. However, consciousness remains unaffected. See the glossary in [Appendix 1](#). The impairments of cognitive function are commonly accompanied, and occasionally preceded, by a deterioration in emotional control, social behaviour, and motivation, and the impairment is sufficient to interfere with everyday activities. Dementia is a collection of different subtypes distinguished by the underlying pathology. ADD is the most common form of dementia and other important pathologies associated with dementia are vascular disease, Lewy bodies, and frontotemporal pathology ([WHO 2012](#)).

Dementia is a serious worldwide public health problem, with a prevalence of 4.7% in adults older than 60 years (6.2% and 6.5% in Europe and the Americas, respectively). Due to its prevalence in older people, it is expected that the number of people with dementia will increase dramatically. Consequently, in the year 2050, an expected number of 115 million people will have dementia. This will result in a considerable economic burden, which currently stands at 1% of the world's Gross National Product (GNP) in direct and indirect costs ([WHO 2012](#)). These financial costs are in addition to the devastating personal and social consequences of the condition.

The definition of MCI applies to people without evidence of significant deterioration in activities of daily living, but with subjective memory complaints and cognitive impairment detected by standardised tests. MCI often precedes clinical dementia, but there is no consensus regarding how to operationalise the MCI diagnosis. There are several clinical criteria to define which people have MCI, including the Petersen criteria or Petersen Revised Criteria ([Petersen 1999](#); [Petersen 2004](#); [Winblad 2004](#)), Clinical Dementia Rating (CDR = 0.5) ([Morris 1993](#)), or 16 other different classifications of MCI ([Matthews 2008](#)).

A diagnosis of MCI reputedly allows testing of preventive interventions that would slow the progression of MCI to dementia. If the progression of MCI to dementia could be deferred by five years, the prevalence of dementia would decrease by 43% in 2050 ([Alzheimer's Association 2010](#)). MCI has an annual progression rate to ADD from 5% to 15%. However, not every person with MCI develops dementia, and a significant number of people recover or stabilise. Therefore, future research should try to clarify which people with MCI develop dementia in order to be able to focus specifically on people who are at high risk of developing dementia. This may possibly explain the failure of therapy to alter the progression to dementia in people with MCI. Other aspects that may contribute to this failure are the disparity in diagnostic criteria and different settings of the studied participants: community, primary, secondary, and research centres ([Bruscoli 2004](#); [Mattsson 2009](#); [Petersen 1999](#); [Petersen 2009](#)).

The definition of Alzheimer's disease pathology is over 100 years old. This pathology includes neuritic plaques that contain deposits of amyloid beta (A β) and neurofibrillary tangles ([Goedert 2006](#)). This pathology is present in approximately 84% of all people with dementia ([Schneider 2007](#)). Furthermore, Alzheimer's disease pathology is found in 88% of people diagnosed with probable

ADD ([Schneider 2009](#)). Despite this, Alzheimer's disease pathology may be found concomitantly at autopsy in people thought to have other forms of dementia, such as vascular dementia, Lewy body dementia, or frontotemporal dementia (FTD) ([Jellinger 2006](#)). Furthermore, at least five common pathologies have been found in the brains of people who died and were thought to have ADD prior to death ([White 2009](#)). Also, Alzheimer's disease pathology was found in 42% of community-dwelling older people without dementia ([Schneider 2007](#)). This has generated controversy about the importance of the presence of Alzheimer's disease pathology. The pathology can be associated with aging per se, and, for older people, the relationship between amyloid plaque burden and cognitive impairment diminishes as age progresses ([Savva 2009](#)). Thus, this pathology could be an epiphenomenon associated with the presence of dementia, e.g. a by-product of repair mechanisms by vascular damage ([De la Torre 2004](#); [Garcia-Alloza 2011](#)). On the other hand, this controversy could be because our clinical diagnostic criteria have not had enough accuracy to diagnose Alzheimer's disease that is detected by histopathology in postmortem studies ([Hyman 2012](#)). In addition, other researchers think that there is not a real controversy about the amyloid hypothesis, because the amyloid cascade and the A β deposition have a primary role in Alzheimer's disease ([Selkoe 2016](#)).

More recently, the development of A β pathology biomarkers in vivo has been suggested as an important advance as a diagnostic tool in the field of Alzheimer's disease, and has promoted the creation of new diagnostic criteria for people without symptoms (preclinical stages), people with MCI, and people with ADD, based on the presence of biomarkers of Alzheimer's disease. These have included A β tracers by positron emission tomography (PET) ([Albert 2011](#); [Dubois 2014](#); [McKhann 2011](#); [Sperling 2011](#)). However, uncertainties regarding the usability of biomarkers in the diagnosis of dementia still exist, mainly due to variation between biomarker types, criteria for positivity, and differences in methodology ([Noel-Storr 2013](#)). This prompted an important initiative, the Standards for Reporting of Diagnostic Accuracy Studies in dementia studies (STARDdem) statement ([Noel-Storr 2014](#)). Consequently, clinical properties of dementia biomarkers should not be assumed, and formal systematic evaluations of sensitivity, specificity, and other properties of biomarkers should be performed ([Davis 2013](#)).

PET is an imaging technique using compounds labelled with short-lived positron-emitting radionuclides. The use of A β ligands permits the in vivo detection of amyloid deposition in the brain. ¹⁸F-florbetaben is a stilbene derivative, which was first described 12 years ago, and is characterised by a high affinity for A β . ¹⁸F-florbetaben has excellent uptake by brain tissue and washout kinetics in mice ([Zhang 2005](#)). ¹⁸F-florbetaben was evaluated in people with ADD, healthy people without ADD ([Barthel 2011](#)), and people with other dementias ([Villemagne 2011](#)).

The Food and Drug Administration (FDA) and the European Medicines Agency (EMA) approved ¹⁸F-florbetaben for A β binding. These agencies have stated that a negative scan indicates sparse or no plaques, which is inconsistent with a diagnosis of ADD, thus effectively excluding this diagnosis. A positive ¹⁸F-florbetaben scan indicates moderate to frequent presence of neuritic amyloid plaques. However, this might also occur in people with other neurological conditions and in older adults with normal cognition. Therefore, it should be combined with other diagnostic evaluations or instruments and cannot be used solely to assess the risk

of progression to ADD. Therefore, a positive result of an ^{18}F -florbetaben scan does not establish the diagnosis of ADD or any other cognitive disorder definitely, and it should be combined with other diagnostic evaluations or instruments. Additionally, the effectiveness and safety of the tests have not been established by predicting development of dementia or other neurological conditions, or by monitoring responses to therapies (EMA 2014; FDA 2014).

Despite not being approved for this purpose by the regulatory agencies, research has been conducted in people with MCI to determine whether biomarkers, such as ^{18}F -florbetaben for A β , increase the risk of developing dementia over time. The evidence for this is uncertain. For this and other reasons, the NIA-AA in the USA established two different criteria for MCI. Firstly, they established the Core Clinical Criteria for use in all clinical settings, without use of biomarkers, and characterised by concerns regarding a change in cognition with impairment in one or more cognitive domains with preservation of independence in functional abilities, therefore no dementia. Secondly, they established the Clinical Research Criteria, which incorporate the use of biomarkers, such as PET amyloid scans, intended for use exclusively in research settings, including academic centres and clinical trials. This will help determine whether positive scans increase the likelihood of progression from MCI to clinical dementia (Albert 2011). Lastly, it is hoped that people with MCI and positive scans will 'enrich' clinical trials, and more people who will progress to dementia in a shorter time will be included to allow more efficient studies of treatments and prevention strategies of ADD (CMS 2013).

An assumption for some researchers, and one on which this systematic review (SR) is predicated, is that if a person has both MCI and the pathology of Alzheimer's disease and develops clinical ADD subsequently, then the cause of the initial MCI and of the ADD was the Alzheimer's pathology. Our approach is an example of assessing diagnostic test accuracy (DTA) using delayed verification of diagnosis. Instead of the reference standard being based on pathology, it is based on a clinical standard and the progression from MCI to ADD, or any other form of non-ADD, or any dementia. Although, for the reasons stated above, a degree of unreliability has been introduced, defining progression has the advantage of being based on what matters most to people with MCI, their families, and clinicians involved in their care.

^{18}F -florbetaben PET scan is considered the diagnostic marker of interest, and in this SR we assessed the DTA of ^{18}F -florbetaben A β binding in the brain and progression of the following:

- From MCI to ADD.
- From MCI to any other form of non-ADD.
- From MCI to any form of dementia

This SR belongs to a series of SRs regarding PET biomarkers for A β , including ^{18}F -florbetapir and ^{18}F -flutemetamol (Martínez 2016).

Target condition being diagnosed

This SR assessed the following three target conditions.

- ADD (progression from MCI to ADD).
- Any other form of dementia (progression from MCI to any other form of non-ADD).

- Any form of dementia (progression from MCI to any form of dementia).

We compared the index test results obtained at baseline with the results of the reference standards obtained at follow-up (delayed verification).

Index test(s)

The ^{18}F -florbetaben scan is an index test for the detection of A β deposition in the brain region of interest (ROI). The ROI is a selected brain area that physicians create for further study in various anatomical areas of the brain. ^{18}F -florbetaben is a molecular biomarker, described as

[^{18}F]BAY 94-9172, trans-4-(N-methyl-amino)-4'-2-[2-(^{18}F)fluoroethoxy]-ethoxy]-ethoxy-stilbene and also referred to as BAY 94-9172 or ZK 6013443, which is a polyethylene glycol stilbene derivative (Zhang 2005).

Image Interpretation

Both the FDA and EMA have described the criteria for ^{18}F -florbetaben A β positivity (EMA 2014; FDA 2014).

^{18}F -florbetaben diagnosis is by PET image assessment, and is defined as positive if the analysis shows the following.

- Moderate or smaller area(s) of tracer uptake equal to or higher than that presented in the white matter: extending beyond the white matter rim to the outer cortical margin involving the majority of the slices within the respective region.
- Pronounced A β deposition (a large confluent area of tracer uptake equal to or higher than that presented in white matter extending beyond the white matter rim to the outer cortical margin and involving the entire region including the majority of slices within the respective region) in the grey matter of the following four brain regions: the temporal lobes, the frontal lobes, the posterior cingulate cortex/precuneus, and the parietal lobes.

Readers trained in PET images with the ^{18}F -florbetaben should interpret the A β PET images made with this ligand (EMA 2014; FDA 2014).

Before the FDA and EMA described the criteria for ^{18}F -florbetaben scan positivity, the diagnosis of dementia was made using different thresholds. Therefore, we planned to use the FDA or EMA criteria applied in each included study to classify participants as either test-positive or test-negative, or, alternatively, if ^{18}F -florbetaben A β uptake and retention exceeded a certain threshold.

We considered the measurement of the ^{18}F -florbetaben retention (retention ratio): distribution volume ratio (DVR), standardised uptake value ratio (SUVR), or other ratios. DVR refers to the ratio of the ^{18}F -florbetaben distribution volume in the selected area (ROI) to the distribution volume in the reference area. SUVR is the ratio of the ^{18}F -florbetaben ligand standardised uptake value in the selected area (ROI) to the standardised uptake value in the reference area.

The unit of analysis of our SR was the participant. We did not include studies that analysed multiple ROIs per person.

Image analysis was not prespecified (e.g. Statistical Parametric Mapping (SPM) or other image analysis techniques).

Administration Instructions and Recommended Dosing

- Time between ^{18}F -florbetaben injection and PET acquisition: images should be acquired in 15 to 20 minutes starting from 45 to 130 minutes after intravenous administration (FDA 2014) or acquired in 20 minutes starting from 90 minutes after intravenous administration (EMA 2014);
- Injection dose: the recommended dose for ^{18}F -florbetaben A β PET is 300 MBq (8.1 mCi), maximum 30 mcg mass dose (FDA 2014) or 300 MBq (240 to 360 MBq) as a single slow intravenous bolus (6 sec/mL) in a total volume of up to 10 mL (EMA 2014).

Although it was inevitable that included studies had used different imaging protocols, readers' expertise, and varied parameters, the amyloid PET data in these included studies should be technically adequate and acquired at a fully qualified and certified facility.

Clinical pathway

At this time, the clinical evaluation often has similarities between different countries (Cordella 2013; NICE 2006). It often starts with people experiencing memory complaints detected by themselves or their relatives. Frequently, general practitioners or family physicians are consulted, and they often conduct a medical evaluation using a screening test for cognitive impairment. Whenever this screening test is positive, they complete an assessment with a clinical evaluation conducted with laboratory studies that can rule out a secondary cause of cognitive impairment (e.g. hypothyroidism, renal failure, liver failure, vitamin B12 or folate deficiency, and others). In addition, these people are then referred to medical specialists in cognitive disorders (preferably a geriatrician, psychiatrist, or neurologist) in a secondary centre or directly to memory clinics where further clinical assessment, laboratory studies, and cerebral image studies are conducted to confirm the dementia diagnosis.

People with dementia, or their relatives, often directly consult these specialists or specialised memory clinics in the study of cognitive disorders. Therefore, the performance of the diagnostic tests will probably vary according to whether it is a primary consultation or referral from primary to specialist care, or if the people have different clinical stages of the disease (MCI, mild, moderate, or severe dementia). Due to these differing pathways, the use of ^{18}F -florbetaben PET ligand for A β is mainly used in specialist consultations and memory clinics as an addition to clinical evaluation or other tests, helping in a clinical setting to discard a diagnosis of Alzheimer's dementia with a negative scan in a person with clinical dementia and doubts about the aetiology (e.g. FTD versus ADD). Otherwise, it might be used solely in the research field in people with MCI for the enrichment of clinical trials, for example, enrolling people with MCI and a positive PET scan to study preventive interventions before people develop dementia.

However, in some memory clinics the ^{18}F -florbetaben PET is used for clinical purposes in people with persistent or progressive unexplained MCI adopting the Johnson criteria (Johnson 2013), criteria without sufficient evidence. Therefore, if the ^{18}F -florbetaben PET is positive in a person with MCI, this positivity is considered as one of the core histopathological findings of Alzheimer's disease. The person will thus be catalogued as a patient

with prodromal Alzheimer's disease or MCI due to Alzheimer's disease.

Alternative test(s)

Currently, there are no standard practice tests available for the clinical diagnosis of Alzheimer's disease dementia. Below, we have listed the alternative tests that we have excluded from this SR. The Cochrane Dementia and Cognitive Improvement Group is in the process of conducting a series of DTA SRs of biomarkers and scales (see list below).

- ^{18}F PET ligands for A β (^{18}F -florbetapir, ^{18}F -flutemetamol) (Martínez 2016).
- ^{18}F -FDG-PET (PET F-fluorodeoxyglucose) (Smailagic 2015).
- ^{11}C -PIB-PET (PET-Pittsburgh compound B) (Zhang 2014).
- Cerebrospinal fluid (CSF) analysis of A β and tau (Kokkinou 2014; Ritchie 2013; Ritchie 2014).
- Structural magnetic resonance imaging (sMRI) (Filippini 2012).
- Neuropsychological tests (Mini-Mental State Examination (MMSE); MiniCOG; Montreal Cognitive Assessment (MoCA) (Arevalo-Rodriguez 2015; Chan 2014; Creavin 2016; Davis 2015; Fage 2015; Seitz 2014).
- Informant interviews (Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE); AD8) (Harrison 2014; Hendry 2014; Lees 2014; Harrison 2015; Quinn 2014).
- APOE- $\epsilon 4$ (Elias-Sonnenschein 2014a; Elias-Sonnenschein 2014b; Elias-Sonnenschein 2014c).
- Single-photon emission computed tomography (SPECT) brain imaging (Archer 2015; McCleery 2015).

Rationale

Accurate and early diagnosis of Alzheimer's disease is crucial for planning in healthcare systems, because the costs of dementia are currently at least 1% of the world's GNP (WHO 2012).

^{18}F -florbetaben is approved for use in the clinical field mainly in people who are diagnosed clinically with dementia of uncertain aetiology, in which case diagnosis of ADD can be discarded if the test is negative. Even though ^{18}F -florbetaben is not approved for this purpose, this biomarker test is currently being used in the research field to search for the accurate identification of people with MCI who would progress to ADD or other forms of dementia. Amyloid β tracers by PET have been included in newly diagnostic criteria in the study in people with MCI (Albert 2011; Dubois 2014). However, some uncertainties exist about the generalisability of the DTA results in clinical settings, especially in older people (Richard 2012).

It is currently believed that if the health system can identify which people are at high risk of progressing from MCI to dementia, it can focus on improving opportunities for appropriate contingency planning for them. Proper recognition of the disease may also help prevent inappropriate and potentially harmful admissions to hospital or institutional care (NAO 2007), and enable the development of new treatments designed to delay or prevent progression to more debilitating stages of the disease. Additionally, this may demonstrate a real clinical benefit for people and caregivers, and will reduce health system costs.

This SR assesses the DTA with ^{18}F -florbetaben A β PET in people with MCI.

OBJECTIVES

To determine the diagnostic test accuracy (DTA) of ^{18}F -florbetaben as the index test for detecting people with mild cognitive impairment (MCI) at time of performing the test who would clinically progress to Alzheimer's disease dementia (ADD), or other forms of non-ADD, or any form of dementia at follow-up.

Secondary objectives

To investigate the heterogeneity of the DTA in the included studies, by evaluating the spectrum of people, referral centres, clinical criteria of MCI, ^{18}F -florbetaben techniques, reference standards used, duration of follow-up, aspects of study quality, and conflicts of interest.

METHODS

Criteria for considering studies for this review

Types of studies

We included longitudinal studies that had prospectively defined cohorts with any accepted definition of mild cognitive impairment (MCI), as outlined below, at time of performing the ^{18}F -florbetaben A β scan and a reference standard (see [Index tests](#) and [Reference standards](#) below). We obtained the results at the follow-up of the studies. These studies had to employ delayed verification of progression to dementia and were sometimes labelled as 'delayed verification cross-sectional studies' ([Bossuyt 2008](#); [Knottnerus 2002](#)). We included case-control studies when they incorporate a delayed verification design. This occurred in the context of a cohort study, so these studies were invariably diagnostic-nested case-control studies.

Participants

Participants recruited and clinically classified as having MCI at time of performing the test were eligible for inclusion. We established the diagnosis of MCI using the Petersen criteria or revised Petersen criteria ([Petersen 1999](#); [Petersen 2004](#); [Winblad 2004](#)), the criteria included in Matthews study ([Matthews 2008](#)), CDR = 0.5 (CDR structured interviews collects information from both the collateral source and the subject regarding memory, orientation, judgment and problem solving, community affairs, home and hobbies, and personal care, where the range of possible scores varies from none = 0 point to severe = 3 points) ([Morris 1993](#)), the National Institute on Aging-Alzheimer's Association (NIA-AA) core clinical criteria ([Albert 2011](#)), or a combination.

We excluded studies that included people with MCI possibly caused by any of the following.

- Current or a history of alcohol or drug abuse.
- Central nervous system (CNS) trauma (e.g. subdural hematoma), tumour, or infection.
- Other neurological conditions (e.g. Parkinson's or Huntington's diseases). Regarding Parkinson's disease, many of the studies specifically excluded people with Parkinson's disease from the group with mild cognitive impairment. This specific group of people is complex in both regards to defining neuropathology

and in determination of functional decline. For these reasons, this group of people needs to be addressed in specific studies.

Index tests

The index test of this SR was ^{18}F -florbetaben biomarker test. We used the criteria and cut-off values for test positivity, as reported in the included studies. We considered positivity for ^{18}F -florbetaben A β scan uptake and retention exceeding a certain threshold.

Target conditions

Three target conditions were included in this SR:

- Alzheimer's disease dementia (ADD) (progression from MCI to ADD).
- Any other forms of dementia (progression from MCI to any other forms of non-ADD).
- Any form of dementia (progression from MCI to any form of dementia).

Reference standards

The reference standard was the progression to the target conditions evaluated by a physician with expertise in the dementia field (preferably a geriatrician, psychiatrist, or neurologist). For the purpose of this SR, we accepted several definitions of ADD. We included studies that applied the National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association (NINCDSADRDA) criteria ([McKhann 1984](#)), the Diagnostic and Statistical Manual of Mental Disorders (DSM) criteria ([APA 1987](#); [APA 1994](#)), and the International Classification of Diseases (ICD) (ICD-10) criteria for ADD. Notably, different iterations of these standards may not be directly comparable over time (e.g. [APA 1987](#) versus [APA 1994](#)). Moreover, the validity of the diagnoses may vary with the degree or manner in which the criteria have been operationalised (e.g. individual clinician versus algorithm versus consensus determination). We considered all these issues when we interpreted the results.

Similarly, we accepted differing clinical definitions of other dementias. For Lewy body dementia, the reference standard is the McKeith criteria ([McKeith 1996](#); [McKeith 2005](#)); for frontotemporal dementia the Lund criteria ([Boxer 2005](#); [Brun 1994](#); [Neary 1998](#)), the DSM criteria ([APA 1987](#); [APA 1994](#)), the ICD criteria (ICD-10), or the International Behavioural Variant FTD Criteria Consortium ([Rascovsky 2011](#)); for vascular dementia, the National Institute of Neurological Disorders and Stroke and Association Internationale pour la Recherche et l'Enseignement en Neurosciences (NINDS-AIREN) criteria ([Román 1993](#)), the DSM criteria ([APA 1987](#); [APA 1994](#)), or the ICD criteria (ICD-10); and, for progressive supranuclear palsy (PSP), the preliminary NINDS criteria ([Hauw 1994](#)).

The time interval over which the progression from MCI to ADD (or other forms of dementia) occurs is very important. We used one year as the minimum period of delay in the verification of the diagnosis (the time between the assessment at which a diagnosis of MCI is made and the assessment at which the diagnosis of dementia is made).

Search methods for identification of studies

Electronic searches

We searched MEDLINE (Ovid SP) from 1946 to May 2017; Embase (Ovid SP) from 1974 to May 2017; PsycINFO (Ovid SP) from 1806 to May 2017; BIOSIS Citation Index (Thomson Reuters Web of Science) from 1922 to May 2017; Web of Science Core Collection, including the Science Citation Index (Thomson Reuters Web of Science) and the Conference Proceedings Citation Index (Thomson Reuters Web of Science) from 1946 to May 2017; LILACS (Bireme); CINAHL (EBSCOhost) from 1980 to May 2017; ClinicalTrials.gov (<https://clinicaltrials.gov>); and the World Health Organization International Clinical Trials Registry Platform (WHO ICTRP) (<http://www.who.int/ictip/search/en/>). We also searched ALOIS, the Cochrane Dementia & Cognitive Improvement Group's specialized register of dementia studies (<http://www.medicine.ox.ac.uk/alosis/>).

We used two approaches in designing the search. One focused solely on the specifically named index test (including a range of synonyms) and the second, run in parallel covered a more general search, linking broader terms for the index test. It focused on terms describing its diagnostic use and terms for the target condition to try to capture the more difficult to locate studies of a more general nature, where these particular radioligands were included in diagnostic accuracy research but not named specifically in the parts of the electronic bibliographic record that are searchable and therefore would be missed.

See [Appendix 2](#) for details of the sources and search strategies that we used. No language or date restrictions were applied to the electronic searches.

Searching other resources

We examined the reference lists of all relevant studies for additional studies. We also searched the Database of Abstracts of Reviews of Effects (DARE) via the Cochrane Library (www.cochranelibrary.com), the National Institute for Health Research - Health Technology Assessment Database (NIHR-HTA) (via the Cochrane Library: www.cochranelibrary.com), the Aggressive Research Intelligence Facility (ARIF) database (www.arif.bham.ac.uk) for other related systematic diagnostic accuracy reviews, and the International Federation of Clinical Chemistry and Laboratory Medicine Committee for Evidence-based Laboratory Medicine database (C-EBLM) (<http://www.ifcc.org/ifcc-education-division/emd-committees/c-ebml/evidence-based-laboratory-medicine-c-ebml-base>).

We checked the reference lists of any relevant studies and SRs, and performed citation tracking using the Science Citation Index to identify any additional relevant studies.

Data collection and analysis

Selection of studies

Two review authors (GM, RV) independently screened the retrieved titles and abstracts for potentially eligible studies. A third review author (PF) resolved any disagreements between the two review authors. The two review authors (GM, RV) then independently assessed the full-text articles of the selected studies with the inclusion criteria. They resolved any disagreements through discussion or, where necessary, consulted a third review author

(PF) who acted as an arbitrator. When a study did not present all relevant data for creating 2 × 2 table, we contacted the study authors directly to request further information. When more than one article presented data on the same population, we included the primary article, which was the article with the largest number of people or with the most informative data (e.g. longest time of follow-up in the primary outcome).

Data extraction and management

We planned to extract the following data regarding the study characteristics.

- Bibliographic details of primary paper:
 - * author, title of study, year, and journal.
- Basic clinical and demographic details:
 - * number of participants;
 - * clinical diagnosis;
 - * MCI clinical criteria;
 - * age;
 - * gender;
 - * sources of referral;
 - * participant recruitment;
 - * sampling procedures.
- Details of the index test:
 - * method of the ¹⁸F-florbetaben administration, including those who administered the test;
 - * thresholds used to define positive and negative tests;
 - * other technical aspects as seemed relevant to the review, e.g. brain areas.
- Details of the reference standard:
 - * definition of ADD and other dementias used in the reference standard;
 - * duration of follow-up from time of the index test performed to defining ADD and other dementias by the reference standard: one year to less than two years; two years to less than four years; and four years or more. If participants had been followed for varied amounts of time, we recorded a mean follow-up period for each included study. If possible, we grouped those data into minimum, maximum, and median follow-up periods, to enable subgroup analyses;
 - * prevalence or proportion of population developing ADD and other dementias, with severity, if described.

We created 2 × 2 tables (cross-relating index test results of the reference standards) as shown in [Appendix 3](#). For the included study, we recorded the number of participants lost to follow-up. We also extracted data necessary for the quality assessment, as defined below. Two review authors (GM, RV) independently performed data extraction. We resolved any disagreements regarding data extraction by discussion, or by consulting a third review author (PF), if it was necessary.

Assessment of methodological quality

We assessed the methodological quality of the included studies using the Quality Assessment of Diagnostic Accuracy Studies 2 tool (QUADAS-2) ([Whiting 2011](#)), as recommended by Cochrane ([Davis 2013](#)). This tool is comprised of four domains: patient selection, index test, reference standard, and patient flow.

Two review authors (GM, RV), who were blinded to each other's scores, independently performed the QUADAS-2 assessment. We resolved any disagreements by discussion or, if necessary, consulted a third review author (PF) who acted as an arbitrator. We assessed each domain in terms of risk of bias, and also considered the first three domains in terms of applicability concerns. In [Appendix 4](#), we have detailed the components of each of these domains and provided a rubric that shows how we made judgements concerning risk of bias. Key areas important to quality assessment were participant selection, blinding, and missing data.

We included three additional signalling questions on our checklist.

- Was the PET scan interpretation done by a trained reader physician? (We included this under the 'Index test' domain.)
- Was there a clear definition of a positive result? (We included this under the 'Index test' domain.)
- Was the study free of commercial funding? (We included this under the 'flow and timing' domain.)

We included the item pertaining to the PET scan interpretation and the definition of positive results to take into account the subjective nature of the ^{18}F -florbetaben A β scan image interpretation, which may be based on a variety of different criteria, such as extensive clinical experience, different standardised uptake values (SUV), different morphological features, or a combination of the aforementioned. We included the third additional item in order to record any potential bias resulting from commercial interest in the results due to the potential risk by the manufacturing company leading to more favourable results and conclusions than sponsorship by other sources ([Lundh 2017](#)).

We did not use QUADAS-2 data to form a summary quality score. We produced a narrative summary that described each included study as at high, low, or unclear risk of bias, as well as concerns regarding applicability, which we have described in [Appendix 5](#).

Statistical analysis and data synthesis

We applied the DTA framework for the analysis of a single test and extracted the data from the study into a 2 x 2 table, showing the binary test results cross-classified with the binary reference standard. We used data from the 2 x 2 tables abstracted from the

included study: true positive (TP), false negative (FN), false positive (FP), true negative (TN), and entered these into Review Manager (RevMan) [Review Manager 2014](#) to calculate the sensitivities, specificities, and their 95% confidence intervals. We also presented the study results graphically by plotting estimates of sensitivities and specificities in a forest plot.

However, due to lack of data, we conducted no meta-analyses. However, we prepared a 'summary of findings table'.

Investigations of heterogeneity

We were able to include only one study, therefore issues of heterogeneity did not arise.

Sensitivity analyses

We found insufficient data to conduct any sensitivity analyses.

Assessment of reporting bias

We did not investigate reporting bias.

RESULTS

Results of the search

The total number of records identified through all databases for this SR was 1382. The PRISMA diagram shows the selection of records through the screening and selection processes ([Figure 1](#)). In total, we assessed 79 studies (42 full text papers, 16 conference publications, 9 registered studies in clinicaltrials.gov, and 12 registered in WHO ICTRP) for eligibility in the full-text screening. We included one study ([Ong 2015](#)). Additionally, five references were identified as ongoing studies (EUCTR2013-004671-12-BE; EUCTR2014-000562-21-NL; EUCTR2014-004244-35-IT; NCT01222351, NCT02854033). We excluded 73 studies: 19 studies were multiple publications or duplicated, and the remaining 53 studies were excluded as they did not meet the inclusion criteria: i) not a longitudinal study (n = 13); ii) not MCI participants at baseline (n = 14); iii) index test not a ^{18}F -florbetaben PET scan (n = 5); iv) discussion or review paper (n = 19); v) wrong outcome or study design (n = 2). One study did not have data suitable for analysis ([Characteristics of excluded studies](#)).

Figure 1. Flow diagram.





Figure 1. (Continued)



Included Study

See [Characteristics of included studies](#).

The study of Ong 2015 was conducted in Australia (Ong 2015). This study included older adult participants who were referred from local memory clinics and who met consensus criteria for MCI at baseline, and were recruited as part of a study to evaluate the ^{18}F -florbetaben PET positivity scan at baseline and progression from MCI to ADD, and compare an SUVR assessment with visually assessed scans in determining a positive or negative scan. The other objective of this study was to examine whether progressive A β accumulation was detectable using the ^{18}F -florbetaben PET scan at follow-up.

Ong 2015 included 45 MCI participants and performed follow-up at two and four years, evaluating progression from MCI to probable ADD. The authors described their ^{18}F -florbetaben status as positive or negative, using a visual assessment by five readers trained on an electronic training tool and their $\text{SUVR} \geq 1.45$ as described previously (Ong 2015). We included the data at four years of follow-up, because, according to the methodology, we included the longest time of follow-up in the primary outcome. MCI participants fulfilled the Petersen 2004 and Winblad 2004 criteria for MCI. Participants had to be at least 60 years old and to have had at least seven years of formal schooling. They were also required to communicate fluently in English, to have no contraindications to undergoing an MRI scan, and to have a MMSE of ≥ 23 points.

There were 21 participants with ^{18}F -florbetaben with an SUVR value < 1.45 and 24 participants with a ^{18}F -florbetaben SUVR value ≥ 1.45 ; using the visual assessment, there were 20 participants with ^{18}F -florbetaben negative and 25 participants with ^{18}F -florbetaben positive. The demographic data provided was based on those classified as positive or negative by SUVR. The age of the participants was 71.8 ± 6.1 and 73.5 ± 6.9 years, years of education 13.5 ± 3.0 , and 13.8 ± 4.2 , and MMSE was 27.9 ± 1.4 and 26.7 ± 1.9 for those with ^{18}F -florbetaben < 1.45 or ≥ 1.45 , respectively. No demographic data was available for those classified by visual assessment. Of the 45 participants classified with SUVR, at four years follow-up, 21 of 45 participants (46.7%) had developed ADD, and 5 of 45 (11.1%) had developed another form of dementia. Of the 45 participants classified with visual assessment, at four years follow-up, 21 of 45 participants (46.7%) had developed ADD, and 5 of 45 (11.1%) had developed another form of dementia. At four years follow-up, the diagnosis was performed by a neurologist with access to all study results and personal medical records. The reference standard was the NINCDS-ADRDA criteria for ADD (McKhann 1984) and, for other forms of dementia, the reference standards were McKeith 1996 for Lewy body dementia, Lund criteria for frontotemporal dementia Neary 1998, and Hauw 1994 for PSP.

Potential conflicts of interest were noted. Financial support for the study was provided by the previous and current manufacturer of the test; three authors were employees from the previous manufacturer of ^{18}F -florbetaben tracer and another three authors were employees from the actual manufacturer of ^{18}F -florbetaben (Ong 2015).

Ongoing studies

Two studies were found as ongoing studies in clinicaltrials.gov. The first study, NCT01222351, included a sub-study of the Washington Heights-Inwood Community Aging Project focused on cognitively normal older adults, older adults with MCI, and older adults with Alzheimer's disease. Participants were selected on the basis of change in plasma amyloid beta levels over prior assessment intervals. The purpose of the study was to examine whether brain amyloid plaque load, which was to be measured with ^{18}F -florbetaben PET scan, varied as a function of change in plasma levels of amyloid beta and the risk and progression of late onset Alzheimer's disease, MCI, and cognitive decline after three years follow-up. No further details were provided regarding index test and reference standard(s). This study has been recruiting participants since December 2010 in the United States. No expected date of publication was provided in this record. The second study, NCT02854033, was focused on cognitively normal, mild cognitive impairment, and mild ADD participants. The main objective was to determine the relationships among the clinical, cognitive, imaging, genetic, and biochemical biomarker characteristics and the evolution of the entire spectrum of ADD and try to identify diagnostic and prognostic markers among others. The clinical follow-up will be five years. No further details were provided regarding the reference standard(s). This study has been recruiting participants since October 2016. No expected date of publication was provided in this record.

Three studies were found as ongoing studies in WHO ICTRP and they belong to the European Union Clinical Trials Register. EUCTR2014-004244-35-IT is a study focused on amnesic MCI participants with long disease duration (range 2 to 10 years) and imaging studies (MRI and/or ^{18}F -FDG-PET) suggestive of involvement of the limbic/mesial temporal lobe, with the main objective to define the value of the load of amyloid protein. The secondary outcome was the correlation of the amyloid load with neuropsychological measures, the clinical indices, the values of A β 42, total tau, and phospho-tau and with data from MRI and ^{18}F -FDG-PET previously acquired for diagnostic purposes and with a clinical follow-up for at least two years in order to assess the possible clinical progression with basal ^{18}F -florbetaben. No further details were provided regarding the participants, index test, and reference standard(s). This study has been ongoing since March 2015. No expected date of publication was provided in this record.

One additional Dutch ongoing study has been found that focuses on an unselected patient population of subjects visiting the memory clinic of the VUmc Alzheimer Center ([EUCTR2014-000562-21-NL](#)). Its main objective was change after ^{18}F -florbetaben in diagnosis, change in level of confidence of diagnosis, and the impact on patient healthcare management, and the secondary outcome in those with MCI was the clinical progression to dementia during annual follow-up (based on follow-up visits to neurologist and neuropsychologist). No further details were provided regarding the participants, index test, and reference standard(s) and the length of follow-up. This study has been ongoing since January 2015. Similarly, no expected date of publication was provided in this record.


One additional Belgian ongoing study has been found that focuses on the predictive value of baseline ^{18}F -florbetaben capture for

longitudinal change in amyloid load measured using PET in MCI cases ([EUCTR2013-004671-12-BE](#)). A secondary outcome was the comparison of CSF A β 42 and amyloid PET for classification of amyloid-positive and amyloid-negative cases, and the comparison of the predictive value of CSF biomarkers A β 42, T-tau and P-tau181P with that of amyloid imaging for MCI cases that progressed to Alzheimer's disease dementia. No further details were provided regarding the index test, and reference standard(s). This study has been ongoing since June 2014 and no expected date of publication has been provided.

Methodological quality of included studies

We assessed methodological quality using the QUADAS-2 tool ([Whiting 2011](#)). Review authors' judgements about each methodological quality item for the included study are presented in the [Characteristics of included studies](#) and in [Figure 2](#).

Figure 2. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each included study



In the patient selection domain, we considered the study of Ong 2015 to be at unclear risk of bias due to lack of reporting on sampling procedures and exclusion criteria ([Ong 2015](#)). We stated that the included study avoided a case-control design because we only considered data on performance of the index test to discriminate between people with MCI who converted to dementia and those who remained stable.

In the index test domain, we considered the study of Ong 2015 at low risk of bias because the positive threshold used in both visual and quantitative assessment was prespecified ([Ong 2015](#)). Moreover, the index test results were interpreted without knowledge of the results of the reference standard. In our two additional signalling questions, the risk concerning the index test being interpreted by a trained reader physician was low, because they were trained on an electronic training tool to do the amyloid PET visual interpretation. The other signalling question was rated as low risk, because there was a clear definition of a positive result.

In the reference standard domain, we considered the study to be at high risk of bias because it was reported that the neurologist had access to all study results and personal medical records to make the diagnosis ([Ong 2015](#)). However, the reference standard(s) were clearly established ([McKhann 1984](#), [McKeith 1996](#) [Neary 1998](#); [Hauw 1994](#)).

In the flow and timing domain, we judged the study to be at high risk of bias because in our additional signalling question there were potential conflicts of interest due to the financial support for the study and, in addition, three authors were employees of the previous manufacturer of ^{18}F -florbetaben tracer and the other three authors were employees of the actual manufacturer of ^{18}F -florbetaben ([Ong 2015](#)). However, the four years of interval between the index test and the reference standard was considered an appropriate interval, all participants received the same reference standard(s), and all 45 participants were accounted for in the analysis.

For assessment of applicability, there was no concern that the included patients and setting, and the conduct and interpretation of the index test, did not match the review question (Ong 2015). However, the target condition (as defined by the reference standard) was of high concern due to the fact that diagnosis was made with full access to study results and medical records at four years follow-up.

Findings

The key characteristics of the study are summarised in [Characteristics of included studies](#). The summary of main results for the only included study is presented in the 'Summary of findings' table ([Summary of findings 1](#)).

Ong 2015 included data on 45 participants with MCI diagnosed with Petersen criteria (Petersen 2004), and Winblad 2004. The study used two different assessments to evaluate the PET: visual assessment

PET positive if increased tracer uptake was visible in any of four different cerebral regions (frontal, parietal, temporal, and posterior cingulate/precuneus cortices) and quantitative assessment with a $SUVR \geq 1.45$ for a positive ^{18}F -florbetaben. At four years follow-up, the diagnosis of ADD was made using NINCDS-ADRDA criteria (McKhann 1984). Lewy Body dementia was made using McKeith 1996 criteria, FTD diagnosis was made using Lund criteria (Neary 1998) and PSP diagnosis was made using (Hauw 1994) criteria.

^{18}F -florbetaben for Alzheimer's disease dementia (ADD)

- **Visual Assessment:** ^{18}F -florbetaben PET scan had a sensitivity of 100% (95% CI 84% to 100%) and a specificity of 83% (95% CI 63% to 95%) to predict the progression from MCI to ADD at four years follow-up. Of 45 participants who were given an initial clinical diagnosis of MCI, 21 were true positive, 4 were false positives, 0 were false negative, and 20 were true negative (Figure 3).

Figure 3. Forest plot of tests: 1 ^{18}F -florbetaben visual assessment and progression to ADD, 2 ^{18}F -florbetaben SUVR and progression to ADD, 3 ^{18}F -florbetaben visual assessment and progression to any other form of non-ADD, 4 ^{18}F -florbetaben SUVR and progression to any other form of non-ADD, 5 ^{18}F -florbetaben visual assessment and progression to any form of dementia, 6 ^{18}F -florbetaben SUVR and progression to any form of dementia.

^{18}F -florbetaben visual assessment and progression to ADD

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Ong 2015	21	4	0	20	1.00 [0.84, 1.00]	0.83 [0.63, 0.95]		

^{18}F -florbetaben SUVR and progression to ADD

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Ong 2015	21	3	0	21	1.00 [0.84, 1.00]	0.88 [0.68, 0.97]		

^{18}F -florbetaben visual assessment and progression to any other form of non-ADD

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Ong 2015	0	25	5	15	0.00 [0.00, 0.52]	0.38 [0.23, 0.54]		

^{18}F -florbetaben SUVR and progression to any other form of non-ADD

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Ong 2015	0	24	5	16	0.00 [0.00, 0.52]	0.40 [0.25, 0.57]		

^{18}F -florbetaben visual assessment and progression to any form of dementia

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Ong 2015	21	4	5	15	0.81 [0.61, 0.93]	0.79 [0.54, 0.94]		

^{18}F -florbetaben SUVR and progression to any form of dementia

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Ong 2015	21	3	5	16	0.81 [0.61, 0.93]	0.84 [0.60, 0.97]		

- **SUVR:** ^{18}F -florbetaben PET scan had a sensitivity of 100% (95% CI 84% to 100%) and a specificity of 88% (95% CI 68% to 97%) to predict the progression from MCI to ADD at four years follow-up. Of 45 participants who were given an initial clinical diagnosis of MCI, 21 were true positive, 3 were false positives, 0 were false negative, and 21 were true negative (Figure 3).

^{18}F -florbetaben for any other form of dementia (non-ADD)

- **Visual Assessment:** ^{18}F -florbetaben PET scan had a sensitivity of 0.0% (95% CI 0.0% to 52%) and a specificity of 38% (95% CI 23% to 54%) to predict the progression from MCI to any other form of dementia (non-ADD) at four years follow-up. Of 45 participants who were given an initial clinical diagnosis of MCI, 0 were true positive; 25 were false positives, 5 were false negative (3 FTD, 1 Lewy body dementia, and 1 PSP), and 15 were true negative (Figure 3).
- **SUVR:** ^{18}F -florbetaben PET scan had a sensitivity of 0.0% (95% CI 0.0% to 52%) and a specificity of 40% (95% CI 25% to 57%) to predict the progression from MCI to any other form of dementia (non-ADD) at four years follow-up. Of 45 participants who were given an initial clinical diagnosis of MCI, 0 were true positive, 24 were false positives, 5 were false negative (3 FTD, 1 Lewy body dementia and 1 PSP), and 16 were true negative (Figure 3).

^{18}F -florbetaben for any form of dementia

- **Visual Assessment:** ^{18}F -florbetaben PET scan had a sensitivity of 81% (95% CI 61% to 93%) and a specificity of 79% (95% CI 54% to 94%) to predict the progression from MCI to any form of dementia at four years follow-up. Of 45 participants who were given an initial clinical diagnosis of MCI; 21 were true positive, 4 were false positives, 5 were false negative (3 FTD, 1 Lewy body dementia, and 1 PSP) and 15 were true negative (Figure 3).
- **SUVR:** ^{18}F -florbetaben PET scan had a sensitivity of 81% (95% CI 61% to 93%) and a specificity of 84% (95% CI 60% to 97%) to predict the progression from MCI to any form of dementia at four years follow-up. Of 45 participants who were given an initial clinical diagnosis of MCI, 21 were true positive, 3 were false positives, 5 were false negative (3 FTD, 1 Lewy body dementia, and 1 PSP), and 16 were true negative (Figure 3).

Investigation of heterogeneity

We were able to include only one study, therefore, issues of heterogeneity did not arise.

Sensitivity analyses

There were insufficient data to permit any sensitivity analyses.

DISCUSSION

Summary of main results

The volume and quality of evidence regarding the DTA of ^{18}F -florbetaben for early diagnosis of ADD and other dementias in people with MCI is very limited. We identified only one study in this systematic review and for that reason we were not able to conduct a meta-analysis, sensitivity analysis, or heterogeneity analyses (Ong 2015). The results are summarised in a 'Summary of findings' table (Summary of findings 1). The study was evaluated as at high risk of bias, mainly due to the potential conflicts of interest

(financial support of the study and also because three authors were employees of the previous company who manufactured the ^{18}F -florbetaben tracer and three authors were employees of the current company that manufactures the ^{18}F -florbetaben tracer (Characteristics of included studies)).

Regarding our objectives: to determine the DTA of the ^{18}F -florbetaben PET scan for detecting people with MCI at baseline who will clinically progress to ADD, or to other forms of dementia or any form of dementia at follow-up, the results were the following:

^{18}F -florbetaben PET scan for Alzheimer's disease dementia (ADD)

Progression from MCI to ADD analysed by visual assessment: sensitivity of 100% (95% CI 84% to 100%) and a specificity of 83% (95% CI 63% to 95%).

Progression from MCI to ADD analysed by $\text{SUVR} \geq 1.45$: sensitivity of 100% (95% CI 84% to 100%) and a specificity of 88% (95% CI 68% to 97%).

^{18}F -florbetaben has a close to perfect sensitivity and a good specificity for predicting progression to ADD through visual assessment evaluation or SUVR (Ong 2015). However, a positive ^{18}F -florbetaben PET scan for $\text{A}\beta$, has been found in other neurological conditions clinically diagnosed, and it was positive in vascular dementia, frontotemporal dementia and dementia with Lewy bodies (Villemagne 2011). Nevertheless, in one study with 12 cases with non-ADD at autopsy, the ^{18}F -florbetaben PET scan was negative in all of them (Sabbagh 2017). On the other hand, in other amyloid biomarkers like PET PiB, the false positive rate could be explained because it has affinity to amyloid in vessel walls, in particular, to cerebral amyloid angiopathy (CAA) (Zhang 2014). We would think that the pathological diagnosis of some patients with clinically probable ADD may be vascular dementia secondary to CAA and some people with MCI may have vascular MCI due to CAA.

As other amyloid tracers, ^{18}F -florbetaben has probed the detection of amyloid plaques that are composed of insoluble $\text{A}\beta$ peptides (EMA 2014, FDA 2014), however, the soluble $\text{A}\beta$ oligomers have been playing a central role in Alzheimer's pathogenesis in the amyloid hypothesis (Heyden 2013), with the possibility of producing false negatives. In addition, amyloid tracers do not bind to the other histopathologic core of Alzheimer's disease, the neurofibrillary tangles (NFTs). There is evidence that indicates that plaques and tangles independently contribute to cognitive impairment over the clinical course of Alzheimer's disease (Serrano-Pozo 2013). Moreover, in another cohort study, the NFT formation might be either unrelated to amyloid plaques formation or a temporally distinct process, or both (Royall 2014). The latest could explain why, in 44 participants with ADD and positive $\text{A}\beta$ histopathology, one had a negative ^{18}F -florbetaben PET scan (Sabri 2015).

Another important factor to be considered in predicting the progression to ADD is the duration of follow-up, because the reported progression rate of MCI to ADD is between 8% and 16% per year (Mitchell 2009). Therefore, a high percentage of people with baseline MCI would progress to Alzheimer's disease dementia if we could include a longer follow-up period, which would consequently affect the predictive accuracy of the ^{18}F -florbetaben PET scan.

However the progression rate at two years was 44% (including two participants who at four years follow-up reverted to MCI or converted to another form of dementia (non-ADD)) and at four years was 47%. This is more than normal and probably can be explained by the setting of recruitment or demographic or MCI characteristics and maybe other underlying factors that can increase the progression rate. In addition, in one systematic review regarding the progression from MCI to ADD with PiB PETp-u, a correlation between longer follow-up and higher specificities was found (Ma 2014). However, due to the lack of data, we were not able to investigate the effect of the follow-up on the progression rate from MCI to ADD, or any form of dementia.

On the other hand, MCI subtypes have been related to progression to dementia. In a large longitudinal study with 550 MCI participants, evidence were found that the MCI subtype, presence of storage memory impairment, multiple domain condition, and presence of APOE $\epsilon 4$ allele increased the risk of progression to dementia. Multivariate survival and Kaplan-Meier analyses showed that amnesic MCI with storage memory impairment had the most and closest risk of progression to dementia (Espinosa 2013). Specifically, in our systematic review, the study of Ong 2015 (Ong 2015), included amnesic and non-amnesic MCI, and after adjusting for both ^{18}F -florbetaben PET scan positive status and hippocampal atrophy, the hazard ratio for the development of ADD from amnesic MCI was not significant. Additionally, some other risk factors like family history of dementia, APOE $\epsilon 4$ allele presence, and A β and tau protein levels in cerebrospinal fluid may contribute to a faster progression rate to dementia. In conclusion, further updated systematic reviews should include high quality research with more detailed data about the characteristics of MCI that are required to not only explore the underlying mechanisms but also to elucidate the causal pathways that link ^{18}F -florbetaben PET scan positivity of diverse MCI subtypes and disease progression.

^{18}F -florbetaben PET scan for any other forms of dementia (non-Alzheimer's disease dementia (non-ADD))

Progression to any other form of non-ADD analysed by visual assessment: sensitivity of 0.0% (95% CI 0.0% to 52%) and a specificity of 38% (95% CI 23% to 54%).

Progression to any other form of non-ADD analysed by $\text{SUVR} \geq 1.45$: sensitivity of 0.0% (95% CI 0.0% to 52%) and a specificity of 40% (95% CI 25% to 57%).

The study reported only 5 people converting to non-ADD at four years follow-up: frontotemporal dementia (3), Lewy body dementia (1) and progressive supranuclear palsy (1); all of them were ^{18}F -florbetaben negative (Ong 2015).

^{18}F -florbetaben PET scan cortical binding has been observed in non-ADD; 9% (1/11) of FTLD, 25% (1/4) of VaD, 29% (2/7) of DLB in a study from Australia (Villemagne 2011) and in 11% (3/27) in those with confirmed non-Alzheimer's disease neurodegenerative pathologies at autopsy (Sabri 2015). However, in this study none of the five converters to non-ADD at four years follow-up were ^{18}F -florbetaben positive; the latest could explain the sensitivity of 0% and specificity of 38% with visual assessment in the included study. Nevertheless, according to histopathological studies, we would expect that studies with more participants, or participants with positive PET scans that progress to non-ADD, could potentially

increase the DTA to predict progression to other forms of dementia and decrease the DTA to predict the progression to ADD.

The latest data suggested the test was insufficient to evaluate the early diagnostic value for progression from MCI to any form of non-ADD.

^{18}F -florbetaben PET scan for any form of dementia

Progression to any form of dementia analysed by visual assessment: sensitivity of 81% (95% CI 61% to 93%) and a specificity of 79% (95% CI 54% to 94%).

Progression to any form of dementia analysed by $\text{SUVR} \geq 1.45$: sensitivity of 81% (95% CI 61% to 93%) and a specificity of 84% (95% CI 60% to 97%).

Ong 2015 reported lower sensitivity and specificity for prediction of any form of dementia other than ADD (Ong 2015). This is explained because the test has a close to perfect sensitivity and good specificity to predict the progression to ADD and if we add the data with those that are PET negative with other types of non-ADD, the sensitivity decreased to 81% and the specificity to 79%.

According to the aforementioned, in the cases of ^{18}F -florbetaben PET scans for any other forms of dementia, there are cases that are PET positive with other neurological conditions in the literature, but not in this study; and this paucity of data with this type of participant, could be explained due to the small sample of participants of this study. For that reason, we would expect that in other studies where we would check the progression to any form of dementia, the DTA would be higher because it is probable to find participants with DBL, FTD and others with a PET positive result, and also we would expect an decrease in the DTA to predict the progression to ADD.

Strengths and weaknesses of the review

We conducted an extensive, comprehensive, and sensitive literature search using 11 different electronic databases without any limitation to language or publication status. However, we only identified one study with 45 eligible participants, therefore our DTA estimates are relatively imprecise. This paucity of evidence reflects the very significant challenges inherent in conducting long term prospective studies of well characterised participants, followed up to the point of progression of a clinical dementia. The methodological quality assessment and data syntheses were based on recommended methods. To increase the reliability of our findings, we included only studies that fulfilled delayed verification of progression from MCI to ADD or any other form of dementia (non-ADD) or any form of dementia at follow-up.

The included study did have significant methodological limitations that weakened confidence in the findings of the review. The study lacked information about the selection of the participants, the reference standard was made with knowledge of the medical studies and medical records, and the major problem was a potential conflict of interest due to the relationship with the companies who produced and produce the tracer. On the other hand, considerable uncertainty remained concerning the clinical diagnosis of ADD; the histopathological diagnosis would be the better way to probe the diagnosis, however, this option is not realistic for a clinical trial.

Applicability of findings to the review question

Regarding the question of this review:

Could the ^{18}F -florbetaben PET scan identify those people with MCI who would progress to clinical dementia at follow-up? There was no applicability concern that the included patients, the setting, the conduct, and interpretation of the index test in the included study did not match the review question. However, there was a high applicability concern about the target condition (as defined by the reference standard) because the diagnosis at follow-up was made with access to the study tests and medical records for all participants and, therefore, due to the one study included, it was difficult to extend the findings into clinical practice without a meta-analysis.

The diagnostic utility of ^{18}F -florbetaben PET scan for identifying Alzheimer's disease pathology and identifying those people with MCI who would convert to ADD could be affected by a number of factors that have not been determined so far. The most important was the lack of a large study to evaluate this question, as we included one study that addressed the question with only 45 participants at follow-up. Conducting a ^{18}F -florbetaben test is expensive, therefore it is important to clearly demonstrate its accuracy prior to recommending its adoption in clinical practice.

AUTHORS' CONCLUSIONS

Implications for practice

Today, the use of ^{18}F -florbetaben is not indicated in people with MCI (FDA and EMA) except in clinical trials and research studies. However, the Amyloid Imaging Task Force, the Society of Nuclear Medicine and Molecular Imaging, and the Alzheimer's Association have proposed the usage of amyloid PET in people with persistent or progressive unexplained MCI (Johnson 2013). The DTA of ^{18}F -florbetaben PET scans, as determined in this SR, suggests a limited use due to a lack of information based only on one study with 45 participants to predict the progression from MCI to ADD and any form of dementia. Despite this, in the sole study, the sensitivity was 100% and the specificity was 83% for visual assessment analysis to predict the progression to ADD and the sensitivity was 81% and the sensitivity was 79% to predict the progression to any form of dementia. The prediction to other forms of dementia (non-ADD) was poor, however, this could be explained because the pathology of the other neurodegenerative conditions is not based on the A β plaques. Finally, we have to consider the risk of bias due to the access to medical tests and medical records by the neurologist who made the diagnosis at four years follow-up, because this could overestimate the DTA of ^{18}F -florbetaben (Lijmer 1999). Due to the aforementioned and the methodological limitations of the included study, it is not possible to recommend the routine use of ^{18}F -florbetaben in clinical practice. The ^{18}F -florbetaben biomarker is expensive, therefore, it is important to clearly demonstrate its DTA and to standardise the process for the diagnostic modality prior to it being widely used.

Implications for research

The FDA and EMA had established the ^{18}F -florbetaben positivity criteria in order that the use in ADD patients' evaluation and use in people with MCI is accepted in research settings and clinical trials (Albert 2011). On the other hand, it has been proposed for

use in clinical practice by the Nuclear Medicine Society and the Alzheimer's Association (Johnson 2013).

The interpretation of the results of the ^{18}F -florbetaben PET scan studies could be difficult due to the use of different methods to define the result of the test. It is still used in many studies with different SUVR, visual assessment or both, and this promotes different accuracies for the tracer, even in people with ADD when they are compared with healthy people without ADD. Therefore, it is necessary to consider that visual assessment is the most important option to interpret the ^{18}F -florbetaben PET scan, because this is the approach to the interpretation established by FDA and EMA (FDA 2014, EMA 2014).

Moreover, clinical assessment in people with memory complaints is not always undertaken with only one test, as clinical assessment could use different tests, like volumetric hippocampal MRI, FDG-PET, SPECT, CSF, and others. This makes sense because neurodegenerative diseases are complex disorders with occasionally multiple and overlapping pathophysiological processes, and multitracer imaging may be helpful in combining metabolic, inflammation, or apoptosis markers with those labelling typical protein aggregations seen in the progression of MCI to Alzheimer's disease dementia. In future, various PET imaging modalities are needed to evaluate the usefulness of the various PET tracers as predictors of progression to Alzheimer's disease dementia in MCI studies with clinical follow-up. There is a hypothesis that amyloid deposition is an early event in Alzheimer's disease that reaches a relative plateau even at the MCI stage, while downstream biomarkers measure neuronal loss and dysfunction, and cognitive measures are more dynamic at the symptomatic disease stage (Jack 2010). Based on this hypothesis, the combination of structural imaging, functional imaging, and cognitive tests may be better predictors of when an individual will convert. However, in this way, the Ong 2015 study showed that a ^{18}F -florbetaben PET positive result predicted progression after adjusting for both aMCI status and hippocampal atrophy evaluated with sMRI (Ong 2015). Nevertheless, there is a lack of studies with ^{18}F -florbetaben combined with other tests.

Additionally, if we consider the hierarchical evidence needed for the level of efficacy of diagnostic imaging tests, we are currently in the second step of six according to Herscovitch (Herscovitch 2015): technical efficacy, diagnostic accuracy efficacy, diagnostic thinking efficacy, therapeutic impact, patient health outcomes, and, finally, societal efficacy. Therefore, we need further research about accuracy before progressing to the other steps with their specific studies before we can incorporate the ^{18}F -florbetaben PET scan into clinical practice.

ACKNOWLEDGEMENTS

Gabriel Martínez is a PhD candidate in the Methodology of Biomedical Research and Public Health at the Department of Paediatrics, Obstetrics and Gynaecology and Preventive Medicine, Universitat Autònoma de Barcelona, Barcelona, Spain.

We are grateful to the authors of the included and excluded studies who responded to our requests for additional information.

We thank the strong support of the Cochrane Dementia and Cognitive Improvement Group (CDCIG), especially Sue Marcus in finalizing the review.

We thank Anna Noel-Storr, Information Specialist of the CDCIG, for her assistance with the design of the search strategy.

We thank Gerard Urrútia and Marta Roqué i Figuls for their contribution in the preparation of the protocol for the review ([Martínez 2016](#))

We thank the peer reviewers for their many helpful suggestions.

REFERENCES

References to studies included in this review

Ong 2015 {published data only}

Bahar-Fuchs A, Villemagne V, Ong K, Chetelat G, Lamb F, Reininger CB, et al. Prediction of amyloid- β pathology in amnesic mild cognitive impairment with neuropsychological tests. *Journal of Alzheimer's Disease* 2013;**33**(2):451-62.

Ong D, Villemagne V, Bahar-Fuchs A, Lamb F, Jones G, Reininger C, et al. Conversion from mild cognitive impairment to Alzheimer's disease over 12 months: predictive value of AB imaging with 18F-Florbetaben. *Internal Medicine Journal* 2011;**41** Suppl S3:39.

Ong K, Villemagne V, Bahar-Fuchs A, Lamb F, Chetelat G, Holl G, et al. Conversion from mild cognitive impairment to Alzheimer's disease over 12 months: predictive value of AB imaging with 18F-florbetaben. *Alzheimer's & Dementia* 2011;**7** Suppl(4):S217.

Ong K, Villemagne V, Bahar-Fuchs A, Lamb F, Reininger C, Putz B, et al. Cognitive change and Ab deposition in MCI subjects studied with serial 18F-florbetaben PET over 2 years. *Alzheimer's & Dementia* 2012;**8** Suppl(4):P22-3.

Ong K, Villemagne VL, Bahar-Fuchs A, Lamb F, Jones G, Reininger C, et al. A two-year longitudinal assessment of abeta deposition in MCI with 18f-florbetaben. *Internal Medicine Journal* 2012;**42** Suppl S3:12.

* Ong KT, Villemagne VL, Bahar-Fuchs A, Lamb F, Langdon N, Catafau AM, et al. A β imaging with 18F-florbetaben in prodromal Alzheimer's disease: a prospective outcome study. *Journal of Neurology, Neurosurgery and Psychiatry* 2015;**86**(4):431-6.

References to studies excluded from this review

Kim 2017 {published data only}

Kim HJ, Seo SW, Kim Y, Jang H, Kim KW, Lee JS, et al. Anatomical subtypes of amnesic mild cognitive impairment. *Neuro-degenerative Diseases* 2017; Vol. 17, issue Suppl 1:1039.

References to ongoing studies

EUCTR2013-004671-12-BE {unpublished data only}

EUCTR2013-004671-12-BE. Predictive value of biomarkers in patients with amnesic mild cognitive impairment. apps.who.int/trialsearch/Trial2.aspx? TrialID=EUCTR2013-004671-12-BE (first received 20 May 2014).

EUCTR2014-000562-21-NL {unpublished data only}

EUCTR2014-000562-21-NL. Amyloid-PET as a diagnostic marker in daily practice. apps.who.int/trialsearch/Trial2.aspx? TrialID=EUCTR2014-000562-21-NL (first received 5 January 2015).

EUCTR2014-004244-35-IT {unpublished data only}

EUCTR2014-004244-35-IT. Amyloid load in prodromal AD with limbic-predominant phenotype [Amyloid load in

prodromal AD with limbic-predominant phenotype principal investigator - Studio del Carico di Amiloide in AD prodromico con fenotipo limbico]. apps.who.int/trialsearch/Trial2.aspx? TrialID=EUCTR2014-004244-35-IT (first received 13 November 2014).

NCT01222351 {unpublished data only}

NCT01222351. Measuring brain amyloid plaque load in older adults using BAY 94-9172. clinicaltrials.gov/show/NCT01222351 (first received 18 October 2010).

NCT02854033 {unpublished data only}

NCT02854033. Alzheimer's disease neuroimaging initiative 3 (ADNI3) protocol. clinicaltrials.gov/show/NCT02854033 (first received 3 August 2016).

Additional references

Albert 2011

Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia* 2011;**7**(3):270-9.

Alzheimer's Association 2010

Alzheimer's Association. Changing the trajectory of Alzheimer's disease: a national imperative. <http://www.alzheimersreadingroom.com/2010/05/changing-trajectory-of-alzheimers.html> (accessed prior to 12 October 2017).

APA 1987

American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 3rd Edition. Washington, DC: American Psychiatric Association, 1987.

APA 1994

American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4th Edition. Washington, DC: American Psychiatric Association, 1994.

Archer 2015

Archer HA, Smailagic N, John C, Holmes RB, Takwoingi Y, Coulthard EJ, et al. Regional Cerebral Blood Flow Single Photon Emission Computed Tomography for detection of Frontotemporal dementia in people with suspected dementia. *Cochrane Database of Systematic Reviews* 2015, Issue 6. [DOI: [10.1002/14651858.CD010896.pub2](https://doi.org/10.1002/14651858.CD010896.pub2)]

Arevalo-Rodriguez 2015

Arevalo-Rodriguez I, Smailagic N, Figuls MRI, Ciapponi A, Sanchez-Perez E, Giannakou A, et al. Mini-Mental State Examination (MMSE) for the detection of Alzheimer's disease and other dementias in people with mild cognitive impairment (MCI). *Cochrane Database of Systematic Reviews* 2015, Issue 3. [DOI: [10.1002/14651858.CD010783.pub2](https://doi.org/10.1002/14651858.CD010783.pub2)]

Barthel 2011

Barthel H, Gertz HJ, Dresel S, Peters O, Bartenstein P, Buerger K, et al. Cerebral amyloid- β PET with florbetaben (18F) in patients with Alzheimer's disease and healthy controls: a multicentre phase 2 diagnostic study. *Lancet. Neurology* 2011;**10**(5):424-35.

Bossuyt 2008

Bossuyt PM, Leeflang MM. Chapter 6: Developing criteria for including studies. In: Deeks JJ, Bossuyt PM, Gatsonis C, editor(s). *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* Version 1.0.0. The Cochrane Collaboration, 2013. Available from srdta.cochrane.org.

Boxer 2005

Boxer AL, Miller BL. Clinical features of frontotemporal dementia. *Alzheimer Disease and Associated Disorders* 2005;**19**(Suppl 1):S3-6.

Brun 1994

Brun A, Englund B, Gustafson L, Passant U, Mann DMA, Neary D, et al. Clinical and neuropathological criteria for frontotemporal dementia. *Journal of Neurology, Neurosurgery and Psychiatry* 1994;**57**(4):416-8.

Bruscoli 2004

Bruscoli M, Lovestone S. Is MCI really just early dementia? A systematic review of conversion studies. *International Psychogeriatrics* 2004;**16**(2):129-40.

Chan 2014

Chan CCH, Fage BA, Smailagic N, Gill SS, Herrmann N, Nikolaou V, et al. Mini-Cog for the diagnosis of Alzheimer's disease dementia and other dementias within a secondary care setting. *Cochrane Database of Systematic Reviews* 2014, Issue 12. [DOI: [10.1002/14651858.CD011414](https://doi.org/10.1002/14651858.CD011414)]

CMS 2013

The Centers for Medicare & Medicaid Services. Decision memo for beta amyloid positron emission tomography in dementia and neurodegenerative disease (CAG-00431N). cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=265 (accessed 08 October 2015).

Cordella 2013

Cordella CB, Borson S, Boustani M, Chodosh J, Reuben D, Verghese J, et al. Alzheimer's Association recommendations for operationalizing the detection of cognitive impairment during the Medicare Annual Wellness Visit in a primary care setting. *Alzheimer's & Dementia* 2013;**9**(2):141-50.

Creavin 2016

Creavin S, Wisniewski S, Noel-Storr A, Trevelyan C, Hampton T, Rayment D, et al. Mini-Mental State Examination (MMSE) for the detection of dementia in clinically unevaluated people aged 65 and over in community and primary care populations. *Cochrane Database of Systematic Reviews* 2016, Issue 4. [DOI: [10.1002/14651858.CD011145.pub2](https://doi.org/10.1002/14651858.CD011145.pub2)]

Davis 2013

Davis DHJ, Creavin ST, Noel-Storr A, Quinn TJ, Smailagic N, Hyde C, et al. Neuropsychological tests for the diagnosis of

Alzheimer's disease dementia and other dementias: a generic protocol for cross-sectional and delayed-verification studies. *Cochrane Database of Systematic Reviews* 2013, Issue 3. [DOI: [10.1002/14651858.CD010460](https://doi.org/10.1002/14651858.CD010460)]

Davis 2015

Davis DHJ, Creavin ST, Yip JLY, Noel-Storr AH, Brayne C, Cullum S. Montreal Cognitive Assessment for the diagnosis of Alzheimer's disease and other dementias. *Cochrane Database of Systematic Reviews* 2015, Issue 10. [DOI: [10.1002/14651858.CD010775.pub2](https://doi.org/10.1002/14651858.CD010775.pub2)]

De la Torre 2004

De la Torre JC. Is Alzheimer's disease a neurodegenerative or a vascular disorder? Data, dogma, and dialectics. *Lancet. Neurology* 2004;**3**(3):184-90.

Dubois 2014

Dubois B, Feldman HH, Jacova C, Hampel H, Molinuevo JL, Blennow K, et al. Advancing research diagnostic criteria for Alzheimer's disease: the IWG-2 criteria. *Lancet. Neurology* 2014;**13**(6):614-29.

Elias-Sonnenschein 2014a

Elias-Sonnenschein LS, Viechtbauer W, Ramakers I, Ukoumunne O, Verhey FRJ, Visser PJ. APOE- ϵ 4 allele for the diagnosis of Alzheimer's and other dementia disorders in people with mild cognitive impairment in a community setting. *Cochrane Database of Systematic Reviews* 2014, Issue 1. [DOI: [10.1002/14651858.CD010948](https://doi.org/10.1002/14651858.CD010948)]

Elias-Sonnenschein 2014b

Elias-Sonnenschein LS, Viechtbauer W, Ramakers I, Ukoumunne O, Verhey FRJ, Visser PJ. APOE- ϵ 4 allele for the diagnosis of Alzheimer's and other dementia disorders in people with mild cognitive impairment in a primary care setting. *Cochrane Database of Systematic Reviews* 2014, Issue 1. [DOI: [10.1002/14651858.CD010949](https://doi.org/10.1002/14651858.CD010949)]

Elias-Sonnenschein 2014c

Elias-Sonnenschein LS, Viechtbauer W, Ramakers I, Ukoumunne O, Verhey FRJ, Visser PJ. APOE- ϵ 4 allele for the diagnosis of Alzheimer's and other dementia disorders in people with mild cognitive impairment in a secondary care setting. *Cochrane Database of Systematic Reviews* 2014, Issue 1. [DOI: [10.1002/14651858.CD010950](https://doi.org/10.1002/14651858.CD010950)]

EMA 2014

European Medicines Agency. Neuraceq. Annex 1. Summary of product characteristics. www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/002553/WC500162592.pdf (accessed 8th April 2015).

Espinosa 2013

Espinosa A, Alegret M, Valero S, Vinyes-Junque G, Hernandez I, Mauleon A, et al. A longitudinal follow-up of 550 mild cognitive impairment patients: evidence for large conversion to dementia rates and detection of major risk factors involved. *Journal of Alzheimer's Disease* 2013;**34**(3):769-80.

Page 2015

Page BA, Chan CCH, Gill SS, Noel-Storr AH, Herrmann N, Smailagic N, et al. Mini-Cog for the diagnosis of Alzheimer's disease dementia and other dementias within a community setting. *Cochrane Database of Systematic Reviews* 2015, Issue 2. [DOI: [10.1002/14651858.CD010860.pub2](https://doi.org/10.1002/14651858.CD010860.pub2)]

FDA 2014

Food, Drug Administration. Neuroceq. www.accessdata.fda.gov/drugsatfda_docs/label/2014/204677s000lbl.pdf (accessed 08/04/2015).

Filippini 2012

Filippini G, Casazza G, Bellatorre AG, Lista C, Duca P, Beecher D, et al. The role of MRI in the early diagnosis of Alzheimer's disease or other dementias in persons with mild cognitive impairment (MCI). *Cochrane Database of Systematic Reviews* 2012, Issue 2. [DOI: [10.1002/14651858.CD009628](https://doi.org/10.1002/14651858.CD009628)]

Garcia-Alloza 2011

Garcia-Alloza M, Gregory J, Kuchibhotla KV, Fine S, Wei Y, Ayata C, et al. Cerebrovascular lesions induce transient β -amyloid deposition. *Brain* 2011;**134**(12):3697-707.

Geslani 2005

Geslani DM, Tierney MC, Herrmann N, Szalai JP. Mild cognitive impairment: an operational definition and its conversion rate to Alzheimer's disease. *Dementia and Geriatric Cognitive Disorders* 2005;**19**(5-6):383-9.

Goedert 2006

Goedert M, Spillantini MG. A century of Alzheimer's disease. *Science* 2006;**314**(5800):777-81.

Harrison 2014

Harrison JK, Fearon P, Noel-Storr AH, McShane R, Stott DJ, Quinn TJ. Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) for the diagnosis of dementia within a general practice (primary care) setting. *Cochrane Database of Systematic Reviews* 2014, Issue 7. [DOI: [10.1002/14651858.CD010771.pub2](https://doi.org/10.1002/14651858.CD010771.pub2)]

Harrison 2015

Harrison JK, Fearon P, Noel-Storr AH, McShane R, Stott DJ, Quinn TJ. Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) for the diagnosis of dementia within a secondary care setting. *Cochrane Database of Systematic Reviews* 2015, Issue 3. [DOI: [10.1002/14651858.CD010772.pub2](https://doi.org/10.1002/14651858.CD010772.pub2)]

Hauw 1994

Hauw JJ, Daniel SE, Dickson D, Horoupian DS, Jellinger K, Lantos PL, et al. Preliminary NINDS neuropathologic criteria for Steele-Richardson-Olszewski syndrome (progressive supranuclear palsy). *Neurology* 1994;**44**(11):2015-9.

Hendry 2014

Hendry K, Lees RA, McShane R, Noel-Storr AH, Stott DJ, Quinn TJ. AD-8 for diagnosis of dementia across a variety of healthcare settings. *Cochrane Database of Systematic Reviews* 2014, Issue 5. [DOI: [10.1002/14651858.CD011121](https://doi.org/10.1002/14651858.CD011121)]

Herscovitch 2015

Herscovitch P. Regulatory approval and insurance reimbursement: the final steps in clinical translation of amyloid brain imaging. *Clinical and Translational Imaging* 2015;**3**:75-7.

Heyden 2013

Hayden EY, Teplow DB. Amyloid β -protein oligomers and Alzheimer's disease. *Alzheimer's Research & Therapy* 2013;**5**(6):1-11.

Hyman 2012

Hyman BT, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Carrillo MC, et al. National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimer's & Dementia* 2012;**8**(1):1-13.

ICD-10

World Health Organization. International statistical Classification of Diseases and related health problems (ICD-10 Version 2010). apps.who.int/classifications/icd10/browse/2010/en (accessed 29 January 2015).

Jack 2010

Jack CR Jr, Knopman DS, Jagust WJ, Shaw LM, Aisen PS, Weiner MW. Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet. Neurology* 2010;**9**(1):119-28.

Jellinger 2006

Jellinger K. Clinicopathological analysis of dementia disorders in the elderly - an update. *Journal of Alzheimer's Disease* 2006;**9**(Supplement 3):61-70.

Johnson 2013

Johnson KA, Minoshima S, Bohnen NI, Donohoe KJ, Foster NL, Herscovitch P, et al. Appropriate use criteria for amyloid PET: a report of the Amyloid Imaging Task Force, the Society of Nuclear Medicine and Molecular Imaging, and the Alzheimer's Association. *Journal of Nuclear Medicine* 2013;**54**(3):476-90.

Knottnerus 2002

Knottnerus JA, Van Weel C, Muris JW. Evaluation of diagnostic procedures. *BMJ* 2002;**324**(7335):477-80.

Kokkinou 2014

Kokkinou M, Smailagic N, Noel-Storr AH, Hyde C, Ukoumunne O, Worrall RE, et al. Plasma and Cerebrospinal fluid (CSF) Abeta42 for the differential diagnosis of Alzheimer's disease dementia in participants diagnosed with any dementia subtype in a specialist care setting. *Cochrane Database of Systematic Reviews* 2014, Issue 1. [DOI: [10.1002/14651858.CD010945](https://doi.org/10.1002/14651858.CD010945)]

Lees 2014

Lees RA, Stott DJ, McShane R, Noel-Storr AH, Quinn TJ. Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) for the early diagnosis of dementia across a variety of healthcare settings. *Cochrane Database of Systematic Reviews* 2014, Issue 10. [DOI: [10.1002/14651858.CD011333](https://doi.org/10.1002/14651858.CD011333)]

Lijmer 1999

Lijmer JG, Mol BW, Heisterkamp S, Bossel GJ, Prins MH, Van der Meulen JH, et al. Empirical evidence of design-related bias in studies of diagnostic tests. *JAMA* 1999;**282**(11):1061-6.

Lundh 2017

Lundh A, Sismondo S, Lexchin J, Busuioac OA, Bero L. Industry sponsorship and research outcome. *Cochrane Database of Systematic Reviews* 2017, Issue 2. [DOI: [10.1002/14651858.MR000033.pub2](https://doi.org/10.1002/14651858.MR000033.pub2)]

Ma 2014

Ma Y, Zhang S, Li J, Zheng DM, Guo Y, Feng J, et al. Predictive accuracy of amyloid imaging for progression from mild cognitive impairment to Alzheimer disease with different lengths of follow-up: a meta-analysis. *Medicine* 2014;**93**(27):1-12.

Martínez 2016

Martínez G, Flicker L, Vernooij RWM, Fuentes Padilla P, Zamora J, Figuls MRI, et al. 18F PET ligands for the early diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI). *Cochrane Database of Systematic Reviews* 2016, Issue 5. [DOI: [10.1002/14651858.CD012216](https://doi.org/10.1002/14651858.CD012216)]

Matthews 2008

Matthews FE, Stephan BC, McKeith IG, Bond J, Brayne C, Medical Research Council Cognitive Function and Ageing Study. Two-year progression from mild cognitive impairment to dementia: to what extent do different definitions agree?. *Journal of the American Geriatrics Society* 2008;**56**(8):1424-33.

Mattsson 2009

Mattsson N, Zetterberg H, Hansson O, Andreasen N, Parnetti L, Jonsson M, et al. CSF biomarkers and incipient Alzheimer disease in patients with mild cognitive impairment. *JAMA* 2009;**302**(4):385-93.

McCleery 2015

McCleery J, Morgan S, Bradley KM, Noel-Storr AH, Ansorge O, Hyde C. Dopamine transporter imaging for the diagnosis of dementia with Lewy bodies. *Cochrane Database of Systematic Reviews* 2015, Issue 1. [DOI: [10.1002/14651858.CD010633.pub2](https://doi.org/10.1002/14651858.CD010633.pub2)]

McKeith 1996

McKeith IG, Galasko D, Kosaka K, Perry EK, Dickson DW, Hansen LA, et al. Consensus guidelines for the clinical and pathologic diagnosis of dementia with Lewy bodies (DLB): report of the consortium on DLB international workshop. *Neurology* 1996;**47**(5):1113-24.

McKeith 2005

McKeith IG, Dickson DW, Lowe J, Emre M, O'Brien JT, Feldman H, et al. Consortium on DLB. Diagnosis and management of dementia with Lewy bodies: third report of the DLB consortium. *Neurology* 2005;**65**(12):1863-72.

McKhann 1984

McKhann GM, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report

of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984;**34**(7):939-44.

McKhann 2011

McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR Jr, Kawas CH, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia* 2011;**7**(3):263-9.

Mitchell 2009

Mitchell AJ, Shiri-Feshki M. Rate of progression of mild cognitive impairment to dementia - meta-analysis of 41 robust inception cohort studies. *Acta Psychiatrica Scandinavica* 2009;**119**(4):252-65.

Morris 1993

Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology* 1993;**43**(11):2412-4.

NAO 2007

National Audit Office. Improving services and support for people with dementia. Report by the Comptroller and Auditor General. HC 604 Session General 2006-2007. www.nao.org.uk/wp-content/uploads/2007/07/0607604.pdf (accessed 25th March 2015).

Neary 1998

Neary D, Snowden JS, Gustafson L, Passant U, Stuss D, Black S, et al. Frontotemporal lobar degeneration: a consensus on clinical diagnostic criteria. *Neurology* 1998;**51**(6):1546-54.

NICE 2006

National Institute for Health Care Excellence. Dementia: supporting people with dementia and their carers in health and social care. NICE guidelines [CG42]. www.nice.org.uk/guidance/cg42 (accessed 17th April 2015).

Noel-Storr 2013

Noel-Storr AH, Flicker L, Ritchie CW, Nguyen GH, Gupta T, Wood P, et al. Systematic review of the body of evidence for the use of biomarkers in the diagnosis of dementia. *Alzheimer's & Dementia* 2013;**9**(3):e96-105.

Noel-Storr 2014

Noel-Storr AH, McCleery JM, Richard E, Ritchie CW, Flicker L, Cullum SJ, et al. Reporting standards for studies of diagnostic test accuracy in dementia: the STARDdem Initiative. *Neurology* 2014;**83**(4):364-73.

Okello 2007

Okello A, Edison P, Archer H, Hinz R, Fox N, Kennedy AM, et al. Amyloid deposition and cerebral glucose metabolism in mild cognitive impairment: a longitudinal 11C-PIB and 18F-FDG PET Study. *Journal of Neurology, Neurosurgery and Psychiatry* 2007;**78**(2):219-20.

Petersen 1999

Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. Mild cognitive impairment: clinical characterization and outcome. *Archives of Neurology* 1999;**56**(3):303-8.

Petersen 2004

Petersen RC. Mild cognitive impairment as a diagnostic entity. *Journal of Internal Medicine* 2004;**256**(3):183-94.

Petersen 2009

Petersen RC, Roberts RO, Knopman DS, Boeve BF, Geda YE, Ivnik RJ, et al. Mild cognitive impairment: ten years later. *Archives of Neurology* 2009;**66**(12):1447-55.

Quinn 2014

Quinn TJ, Fearon P, Noel-Storr AH, Young C, McShane R, Stott DJ. Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) for the diagnosis of dementia within community dwelling populations. *Cochrane Database of Systematic Reviews* 2014, Issue 4. [DOI: [10.1002/14651858.CD010079.pub2](https://doi.org/10.1002/14651858.CD010079.pub2)]

Rascovsky 2011

Rascovsky K, Hodges JR, Knopman D, Mendez MF, Kramer JH, Neuhaus J, et al. Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain* 2011;**134**(Pt 9):2456-77.

Review Manager 2014 [Computer program]

Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Richard 2012

Richard E, Schmand B, Eikelenboom P, Westendorp RG, Van Gool WA. The Alzheimer myth and biomarker research in dementia. *Journal of Alzheimer's Disease* 2012;**31**(Suppl 3):S203-9.

Ritchie 2013

Ritchie C, Smailagic N, Ladds EC, Noel-Storr AH, Ukoumunne O, Martin S. CSF tau and the CSF tau/ABeta ratio for the diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI). *Cochrane Database of Systematic Reviews* 2013, Issue 10. [DOI: [10.1002/14651858.CD010803](https://doi.org/10.1002/14651858.CD010803)]

Ritchie 2014

Ritchie C, Smailagic N, Noel-Storr AH, Takwoingi Y, Flicker L, Mason SE, et al. Plasma and cerebrospinal fluid amyloid beta for the diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI). *Cochrane Database of Systematic Reviews* 2014, Issue 6. [DOI: [10.1002/14651858.CD008782.pub4](https://doi.org/10.1002/14651858.CD008782.pub4)]

Román 1993

Román GC, Tatemichi TK, Erkinjuntti T, Cummings JL, Masdeu JC, Garcia JH, et al. Vascular dementia: diagnostic criteria for research studies. Report of the NINDS-AIREN International Workshop. *Neurology* 1993;**43**(2):250-60.

Royall 2014

Royall DR, Palmer RF. The temporospatial evolution of neuritic plaque-related and independent tauopathies: implications for dementia staging. *Journal of Alzheimer's Disease* 2014;**40**(3):541-9.

Sabbagh 2017

Sabbagh MN, Schäuble B, Anand K, Richards D, Murayama S, Akatsu H, et al. Histopathology and florbetaben PET in patients incorrectly diagnosed with Alzheimer's disease. *Journal of Alzheimer's Disease* 2017;**56**(2):441-46.

Sabri 2015

Sabri O, Sabbagh MN, Seibyl J, Barthel H, Akatsu H, Ouchi Y, et al. Florbetaben PET imaging to detect amyloid beta plaques in Alzheimer's disease: phase 3 study. *Alzheimer's & Dementia* 2015;**11**(8):964-74.

Savva 2009

Savva GM, Wharton SB, Ince PG, Forster G, Matthews FE, Brayne C, Medical Research Council Cognitive Function and Ageing Study. Age, neuropathology, and dementia. *New England Journal of Medicine* 2009;**360**(22):2302-9.

Schneider 2007

Schneider JA, Arvanitakis Z, Bang W, Bennett DA. Mixed brain pathologies account for most dementia cases in community-dwelling older persons. *Neurology* 2007;**69**(24):2197-204.

Schneider 2009

Schneider JA, Arvanitakis Z, Leurgans SE, Bennett DA. The neuropathology of probable Alzheimer disease and mild cognitive impairment. *Annals of Neurology* 2009;**66**(2):200-8.

Seitz 2014

Seitz DP, Fage BA, Chan CCH, Gill SS, Herrmann N, Smailagic N, et al. Mini-Cog for the diagnosis of Alzheimer's disease dementia and other dementias within a primary care setting. *Cochrane Database of Systematic Reviews* 2014, Issue 12. [DOI: [10.1002/14651858.CD011415](https://doi.org/10.1002/14651858.CD011415)]

Selkoe 2016

Selkoe DJ, Hardy J. The amyloid hypothesis of Alzheimer's disease at 25 years. *European Molecular Biology Organization* 2016;**8**(6):595-608.

Serrano-Pozo 2013

Serrano-Pozo A, Qian J, Monsell SE, Frosch MP, Betensky RA, Hyman BT. Examination of the clinicopathologic continuum of Alzheimer disease in the autopsy cohort of the National Alzheimer Coordinating Center. *Journal of Neuropathology and Experimental Neurology* 2013;**72**(12):1182-92.

Smailagic 2015

Smailagic N, Vacante M, Hyde C, Martin S, Ukoumunne O, Sachpekidis C. ¹⁸F-FDG PET for the early diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI). *Cochrane Database of Systematic Reviews* 2015, Issue 1. [DOI: [10.1002/14651858.CD010632.pub2](https://doi.org/10.1002/14651858.CD010632.pub2)]

Sperling 2011

Sperling RA, Aisen PS, Beckett LA, Bennett DA, Craft S, Fagan AM, et al. Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia* 2011;**7**(3):280-92.

Villemagne 2011

Villemagne VL, Ong K, Mulligan RS, Holl G, Pejoska S, Jones G, et al. Amyloid imaging with (18)F-florbetaben in Alzheimer disease and other dementias. *Journal of Nuclear Medicine* 2011;**52**(8):1210-7.

Visser 2006

Visser PJ, Kester A, Jolles J, Verhey F. Ten-year risk of dementia in subjects with mild cognitive impairment. *Neurology* 2006;**67**(7):1201-7.

White 2009

White L. Brain lesions at autopsy in older Japanese-American men as related to cognitive impairment and dementia in the final years of life: a summary report from the Honolulu-Asia aging study. *Journal of Alzheimer's Disease* 2009;**18**(3):713-25.

Whiting 2011

Whiting PF, Rutjes AWS, Westwood ME, Mallet S, Deeks JJ, Reitsma JB. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Annals of Internal Medicine* 2011;**155**(8):529-36.

WHO 2012

World Health Organization, Alzheimer's Disease International. Dementia: a public health priority. www.who.int/mental_health/publications/dementia_report_2012/en/ (accessed 23th September 2015).

Winblad 2004

Winblad B, Palmer K, Kivipelto M, Jelic V, Fratiglioni L, Wahlund LO, et al. Mild cognitive impairment - beyond controversies, towards a consensus: report of the International Working Group on Mild Cognitive Impairment. *Journal of Internal Medicine* 2004;**256**(3):240-6.

Zhang 2005

Zhang W, Oya S, Kung MP, Hou C, Maier DL, Kung HF. F-18 Polyethyleneglycol stilbenes as PET imaging agents targeting Abeta aggregates in the brain. *Nuclear Medicine and Biology* 2005;**32**(8):799-809.

Zhang 2014

Zhang S, Smailagic N, Hyde C, Noel-Storr AH, Takwoingi Y, McShane R, et al. 11C-PIB-PET for the early diagnosis of Alzheimer's disease dementia and other dementias in people with mild cognitive impairment (MCI). *Cochrane Database of Systematic Reviews* 2014, Issue 7. [DOI: [10.1002/14651858.CD010386.pub2](https://doi.org/10.1002/14651858.CD010386.pub2)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Ong 2015

Study characteristics	
Patient sampling	<p>Forty-five older adults who were referred from local memory clinics in Australia. No further details of patient sampling and recruitment were reported.</p> <p>Participants had to be at least 60 years old and to have had at least 7 years of formal schooling. They were also required to communicate fluently in English.</p> <p>Exclusion criteria included the presence of dementia, a score lower than 23 on the MMSE, the presence of other conditions that may impair their cognition and independence, including other neurological (stroke, multiple sclerosis, epilepsy, moderate-severe traumatic brain injury), psychiatric (psychotic symptoms, bipolar disorder), or substance use conditions (e.g. drug and alcohol dependence, use of acetylcholinesterase inhibitors and memantine).</p>
Patient characteristics and setting	<p>45 MCI participants diagnosed by the Petersen 2004 and Winblad 2004 criteria.</p> <p>Demographic data were reported for 45 MCI participants (with SUVR); demographic data were not available for those classified by visual assessment.</p> <p>¹⁸F-florbetaben positive: 24, ¹⁸F-florbetaben negative: 21</p> <p>Gender: not described</p> <p>Age Mean (SD): ¹⁸F-florbetaben positive: 73.5 (6.9), ¹⁸F-florbetaben negative: 71.8 (6.1) years</p>

Ong 2015 (Continued)

APOE ε4 carrier: not reported

MMSE Mean (SD) ¹⁸F-florbetaben positive: 26.7 (1.9), ¹⁸F-florbetaben negative: 27.9 (1.4)

Years of education Mean (SD): ¹⁸F-florbetaben positive: 13.8 (4.2), ¹⁸F-florbetaben negative: 13.5 (3.0)

Sources of referral: not reported

Setting: secondary care (memory clinic)

Index tests

Administration instructions and tracer dosis

¹⁸F-florbetaben was injected intravenously over 38 ± 17 s, with a mean specific activity at the time of injection of 60 ± 29 GBq/μmol. Each participant received on average 286 ± 19 MBq of ¹⁸F-florbetaben.

PET imaging was conducted using a 3D GSO Phillips Allegro PET camera. A 2-min transmission scan using a rotating ¹³⁷Cs source was done for attenuation correction immediately prior to scanning. Images obtained between 90 to 110 min post injection were analysed. Images were reconstructed using a 3D RAMLA.

Image interpretation

PET images were processed with a semiautomatic volume of interest (VOI) method. This method used a preset template of narrow cortical VOI that was applied to either the spatially normalized ¹⁸F-florbetaben scan or via placement on the subject's spatially normalized coregistered MRI by a single operator who was blinded to the subject's clinical status. Minor manual adjustments on the MRI were made to ensure that overlap with white matter and cerebrospinal fluid was minimized.

Spatial normalization and coregistration of the PET and MRI images was performed using SPM8.

Mean radioactivity values were obtained from VOI for cortical, subcortical, and cerebellar regions. The cerebellar cortical VOI was placed taking care to avoid cerebellar white matter. No correction for partial volume effect was applied to the PET data.

- **Visual assessment**

Baseline ¹⁸F-florbetaben PET images underwent visual assessment by five independent nuclear medicine physicians blinded to clinical data. Readers had limited or no prior experience with amyloid PET visual interpretation and were trained on an electronic training tool.

The image assessment was performed on axial slices, in a grey scale. The regional cortical tracer uptake (RCTU) scoring system was used to assess the beta-amyloid deposition in the following four regions: frontal cortex, posterior cingulate/precuneus, lateral temporal cortex, and parietal cortex. A RCTU value of 1, 2, or 3 was assigned if either no, moderate, or pronounced tracer uptake was observed respectively as described below:

1 No tracer uptake: Tracer uptake (i.e. signal intensity) in grey matter in the region is lower than in white matter

2 Moderate tracer uptake: Smaller area(s) of tracer uptake equal to or higher than that present in white matter;

- extending beyond the white matter rim to the outer cortical margin.

- involving the majority of the slices within the respective region.

3 Pronounced tracer uptake: A large confluent area of tracer uptake equal to or higher than that present in white matter;

- extending beyond the white matter rim to the outer cortical margin.

- and involving the entire region including the majority of the slices within the respective region.

The RCTU scores for the four brain regions were condensed into a binary brain amyloid plaque load (BAPL) score with positive or negative interpretation as described below:

Ong 2015 (Continued)

¹⁸F-florbetaben PET scan negative: scan without β -amyloid deposition (RCTU score 1 in each of the 4 brain regions 1, 2, 3, and 4).

¹⁸F-florbetaben PET scan positive: scan with moderate β -amyloid deposition (RCTU score 2 in any or all of the 4 brain regions 1, 2, 3, and 4 and no score 3 in these 4 regions) or scan with pronounced β -amyloid deposition (RCTU score 3 at least in one of the brain regions 1, 2, 3, and 4).

The majority read approach established the final result.

• **Quantitative assessment (SUVR):**

PET images were processed with a semiautomatic volume of interest (VOI) method. This method used a preset template of narrow cortical VOI that was applied to either the spatially normalized ¹⁸F-florbetaben scan or via placement on the subject's spatially normalised coregistered MRI by a single operator who was blinded to the subject's clinical status. Minor manual adjustments on the MRI were made to ensure that overlap with white matter and cerebrospinal fluid was minimized. Spatial normalization and coregistration of the PET and MRI images was performed using SPM8. Mean radioactivity values were obtained from VOI for cortical, subcortical, and cerebellar regions. The cerebellar cortical VOI were placed taking care to avoid cerebellar white matter. No correction for partial volume effect was applied to the PET data.

The standardised uptake value (SUV), defined as the decay-corrected brain radioactivity concentration normalised for injected dose and body weight, was calculated for all regions. These were then used to derive the SUV ratio (SUVR), which was referenced to cerebellar cortex. Neocortical A β burden was expressed as the average SUVR of the area-weighted mean for the following cortical ROIs: frontal (consisting of dorsolateral prefrontal, ventrolateral prefrontal, and orbitofrontal regions), superior parietal, lateral temporal, lateral occipital, and anterior and posterior cingulate. The prespecified SUVR used was ≥ 1.45 .

Target condition and reference standard(s)	Target condition as ADD with NINCDS-ADRDA criteria were established with access to all study results and personal medical records
Flow and timing	<p>Duration of follow-up: 4 years</p> <p>Number included in analysis: 45 participants</p> <p>Visual Assessment: 25 ¹⁸F-florbetaben (+) and 20 ¹⁸F-florbetaben (-)</p> <p>Progression from MCI to ADD by visual assessment:</p> <p>¹⁸F-florbetaben (+): 21 MCI to ADD and 4 MCI-MCI; ¹⁸F-florbetaben (-): 0 MCI to ADD and 20 MCI-MCI</p> <p>TP = 21; FP = 4; FN = 0; TN = 20</p> <p>Progression from MCI to any other form of dementia (non-ADD) by visual assessment:</p> <p>¹⁸F-florbetaben (+): 0 MCI to non-ADD and 25 MCI-MCI; ¹⁸F-florbetaben (-): 5 MCI to non-ADD and 15 MCI-MCI</p> <p>TP = 0; FP = 25; FN = 5; TN = 15</p> <p>Progression from MCI to any form of dementia by visual assessment:</p> <p>¹⁸F-florbetaben (+): 21 MCI to any dementia and 4 MCI-MCI; ¹⁸F-florbetaben (-): 5 MCI to any dementia and 15 MCI-MCI</p> <p>TP = 21; FP = 4; FN = 5; TN = 15</p> <p>SUVR: 24 ¹⁸F-florbetaben (+) and 21 ¹⁸F-florbetaben (-)</p> <p>Progression from MCI to ADD by SUVR:</p> <p>¹⁸F-florbetaben (+): 21 MCI to ADD and 3 MCI-MCI; ¹⁸F-florbetaben (-): 0 MCI to ADD and 21 MCI-MCI</p> <p>TP = 21; FP = 3; FN = 0; TN = 21</p>

Ong 2015 (Continued)

Progression from MCI to any other form of dementia (non-ADD) by SUVR:
¹⁸F-florbetaben (+): 0 MCI to non-ADD and 24 MCI-MCI or ADD; ¹⁸F-florbetaben (-): 5 MCI to non-ADD and 16 MCI-MCI or ADD

TP = 0; FP = 24; FN = 5; TN = 16

Progression from MCI to any form of dementia by SUVR:
¹⁸F-florbetaben (+): 21 MCI to any dementia and 3 MCI-MCI; ¹⁸F-florbetaben (-): 5 MCI to any dementia and 16 MCI-MCI

TP = 21; FP = 3; FN = 5; TN = 16

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Unclear		
		Unclear	Low
DOMAIN 2: Index Test All tests			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
Was the PET scan interpretation done by a trained reader physician?	Yes		
Was there a clear definition of a positive result?	Yes		

Ong 2015 (Continued)

	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	No	
	High	High
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	Yes	
Was the study free of commercial funding?	No	
	High	

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Kim 2017	No data for constructing a 2 x 2 table was provided. The study focused on different cortical thickness subgroups and the progression to dementia in aMCI participants whom were evaluated at baseline with PiB PET or ¹⁸ F-florbetaben. We emailed the authors to resolve this issue, however, no response from the lead author was received.

ADD: Alzheimer's disease dementia

APOE: Apolipoprotein E

BAPL: Brain amyloid plaque load

FN: False negative

FP: False positive

MCI: Mild cognitive impairment

MMSE: Mini Mental State Examination

MRI: Magnetic resonance imaging

NINCDS-ADRDA: National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association

PET: Positron emission tomography

RCTU: Regional cortical tracer uptake

RAMLA: Row action maximum likelihood algorithm

ROI: Region of interest

SD: Standard deviation

SPM8: Statistical parametric mapping 8

SUV: Standardised uptake value

SUVR: Standardised uptake value ratio

TN: True negative

TP: True positive

VOI: Volume of interest

Characteristics of ongoing studies *[ordered by study ID]*

EUCTR2013-004671-12-BE

Trial name or title	Predictive value of biomarkers in patients with amnesic mild cognitive impairment
Target condition and reference standard(s)	Cognitive decline after two years of follow-up, reference standard not specified
Index and comparator tests	¹⁸ F-florbetaben
Starting date	June 2014
Contact information	rik.vandenberghe@uzleuven.be University of Leuven
Notes	Dr. Vandenberghe was contacted; he provided requested information regarding the tracer used; email from Dr. Vandenberghe on 23/01/17

EUCTR2014-000562-21-NL

Trial name or title	Amyloid-PET as a diagnostic marker in daily practice
Target condition and reference standard(s)	Progression to dementia, reference standard not specified
Index and comparator tests	¹⁸ F-florbetaben
Starting date	January 2015
Contact information	Alzheimer Center, VU University Medical Center Alzheimer Center alzheimercentrum@vumc.nl
Notes	

EUCTR2014-004244-35-IT

Trial name or title	Amyloid load in prodromal ADD with limbic-predominant phenotype
Target condition and reference standard(s)	Clinical 'indices', reference standard not specified
Index and comparator tests	¹⁸ F-florbetaben
Starting date	March 2015
Contact information	Medicina Nucleare, IRCCS Ospedale San Raffaele perani.daniela@hsr.it
Notes	

NCT01222351

Trial name or title	BAY 94-9172 PET/CT in cognitively normal older adults, older adults with mild cognitive impairment, and older adults with Alzheimer's disease
Target condition and reference standard(s)	Alzheimer's disease, reference standard not specified
Index and comparator tests	¹⁸ F-florbetaben
Starting date	2010
Contact information	Oksana Taterina, ot2004@columbia.edu Yaakov Stern, ys11@columbia.edu
Notes	

NCT02854033

Trial name or title	Alzheimer's disease neuroimaging initiative 3 (ADNI3) protocol
Target condition and reference standard(s)	Rate of progression to dementia due to ADD, reference standard not specified
Index and comparator tests	¹⁸ F-florbetaben, ¹⁸ F-florbetapir
Starting date	October 2016
Contact information	Dr. Paul Aisen, Director, Alzheimer's Therapeutic Research Institute, University of Southern California
Notes	

ADD: Alzheimer's disease dementia

ADNI3: Alzheimer's disease neuroimaging initiative 3

CT: Computed tomography

PET: Positron emission tomography

DATA

Presented below are all the data for all of the tests entered into the review.

Table Tests. Data tables by test

Test	No. of studies	No. of participants
1 ¹⁸ F-florbetaben visual assessment and progression to ADD	1	45
2 ¹⁸ F-florbetaben SUVR and progression to ADD	1	45
3 ¹⁸ F-florbetaben visual assessment and progression to any other form of non-ADD	1	45
4 ¹⁸ F-florbetaben SUVR and progression to any other form of non-ADD	1	45
5 ¹⁸ F-florbetaben visual assessment and progression to any form of dementia	1	45
6 ¹⁸ F-florbetaben SUVR and progression to any form of dementia	1	45

Test 1. ¹⁸F-florbetaben visual assessment and progression to ADD.

Test 2. ¹⁸F-florbetaben SUVR and progression to ADD.

Test 3. ¹⁸F-florbetaben visual assessment and progression to any other form of non-ADD.

Test 4. ¹⁸F-florbetaben SUVR and progression to any other form of non-ADD.

Test 5. ¹⁸F-florbetaben visual assessment and progression to any form of dementia.

Test 6. ¹⁸F-florbetaben SUVR and progression to any form of dementia.

APPENDICES

Appendix 1. Glossary

Aetiology: the cause, set of causes, or manner of causation of a disease or condition.

Amyloid beta (Aβ): an amyloid that is derived from a larger precursor protein and is the primary component of plaques characteristic of Alzheimer's disease.

Biomarker: measurable and quantifiable biological parameters (e.g., specific enzyme concentration, specific hormone concentration, presence of biological substances) which serve as indices for health- and physiology-related assessments, such as disease risk, psychiatric disorders, environmental exposure and its effects, disease diagnosis; metabolic processes; etc.

Bolus: a single dose of a drug or other medicinal preparation given all at once.

Cingulate cortex: one of the convolutions on the medial surface of the cerebral hemispheres.

Cortical: the thin layer of grey matter on the surface of the cerebral hemispheres. It reaches its highest development in humans and is responsible for intellectual faculties and higher mental functions.

Epiphenomenon: A secondary effect or by-product. A secondary symptom or pathology, occurring simultaneously with a disease or condition but not directly related to it.

Frontotemporal: relating to the frontal and the temporal cerebral lobes.

Histopathology: the study of changes in tissues caused by disease.

Hypothyroidism: a syndrome that results from abnormally low secretion of thyroid hormones from the thyroid gland.

Index test: the test under evaluation.

In vivo: (of processes) performed or taking place in a living organism.

Ligand: a molecule that binds to another molecule, used especially to refer to a small molecule that binds specifically to a larger molecule, e.g., an antigen binding to an antibody, a hormone or neurotransmitter binding to a receptor, or a substrate or allosteric effector binding to an enzyme.

Neuritic plaques: accumulations of extracellularly deposited amyloid fibrils within tissues. Is one of the hallmarks of Alzheimer's disease.

Neurofibrillary tangles: abnormal structures located in various parts of the brain and composed of dense arrays of paired helical filaments (neurofilaments and microtubules). Are aggregates of hyperphosphorylated tau protein that are most commonly known as a primary marker of Alzheimer's disease.

Parietal lobe: upper central part of the cerebral hemisphere. It is located anterior to the occipital lobe, and superior to the temporal lobes.

Positron: an extremely small piece of matter with a positive electrical charge, having the same mass as an electron.

Precuneus: is a part of the parietal lobe of the brain, lying on the medial surface of the cerebral hemisphere.

Prodromal: Relating to prodrome; indicating an early stage of a disease.

Radionuclide (sometimes called a radioisotope or isotope): is a chemical which emits a type of radioactivity called gamma rays. The radioactivity can be detected by special scanners.

Reference standard: the best available method for establishing the presence or absence of the target condition.

Sensitivity: a measure of a test's ability to correctly detect people with the disease. It is the proportion of diseased cases that are correctly identified by the test. It is calculated as follows: Sensitivity = Number with disease who have a positive test/Number with disease.

Specificity: a measure of a test's ability to correctly identify people who do not have the disease. It is the proportion of people without the target disease who are correctly identified by the test. It is calculated as follows: Specificity = Number without disease who have a negative test/Number without disease.

Stilbene: organic compounds that contain 1,2-diphenylethylene as a functional group.

Target condition: the disease or condition that the index test is expected to detect.

Temporal lobe: lower lateral part of the cerebral hemisphere responsible for auditory, olfactory, and semantic processing. It is located inferior to the lateral fissure and anterior to the occipital lobe.

Vascular: relating to, affecting, or consisting of a vessel or vessels, especially those which carry blood.

Appendix 2. Search strategy for ¹⁸F-florbetaben PET ligand

Source	Search strategy
MEDLINE In-process and other non-indexed citations and Medline® 1946 to May 2017 (Ovid SP)	<ol style="list-style-type: none"> 1. Florbetaben.ti,ab,nm. 2. (NEURACEQ or neuraceq*).ti,ab,nm. 3. "florbetaben-fluorine-18".ti,ab,nm. 4. "18F-BAY94-9172".ti,ab,nm. 5. "[18F]Florbetaben".ti,ab,nm. 6. "florbetaben-PET".ti,ab,nm. 7. or/1-6 8. Fluorine Radioisotopes/du 9. Aniline Compounds/du 10. Ethylene Glycols/du 11. Stilbenes/du 12. Radioligand Assay/ 13. radioligand*.ti,ab. 14. or/8-13 15. Alzheimer Disease/ri [Radionuclide Imaging] 16. Plaque, Amyloid/ri [Radionuclide Imaging] 17. or/15-16 18. 14 and 17 19. 7 or 18
Embase 1974 to May 2017 (Ovid SP)	<ol style="list-style-type: none"> 1. Florbetaben.ti,ab. 2. (NEURACEQ or neuraceq*).ti,ab. 3. "florbetaben-fluorine-18".ti,ab. 4. "18F-BAY94-9172".ti,ab. 5. "[18F]Florbetaben".ti,ab. 6. "florbetaben-PET".ti,ab. 7. exp florbetaben f 18/ 8. or/1-7 9. exp *radioligand/ 10. Alzheimer disease/ 11. Alzheimer*.ti,ab. 12. amyloid plaque/di [Diagnosis] 13. mild cognitive impairment/ 14. or/10-13 15. 9 and 14 16. 8 or 15
PsycINFO 1806 to May 2017 (Ovid SP)	<ol style="list-style-type: none"> 1. Florbetaben.ti,ab. 2. (NEURACEQ or neuraceq*).ti,ab. 3. "florbetaben-fluorine-18".ti,ab.

(Continued)

	4. "18F-BAY94-9172".ti,ab. 5. "[18F]Florbetaben".ti,ab. 6. "florbetaben-PET".ti,ab. 7. or/1-6
BIOSIS Citation Index (Thomson Reuters Web of Science) (1922 to May 2017)	Topic=(Florbetaben OR NEURACEQ OR neuraceq* OR "florbetaben-fluorine-18" OR "18F-BAY94-9172" OR "[18F]Florbetaben" OR "florbetaben-PET") Timespan=All years. Databases=BCI
Web of Science Core Collection, including the Science Citation Index and the Conference Proceedings Citation Index (Thomson Reuters Web of Science) (1946 to May 2017)	Topic=(Florbetaben OR NEURACEQ OR neuraceq* OR "florbetaben-fluorine-18" OR "18F-BAY94-9172" OR "[18F]Florbetaben" OR "florbetaben-PET") Timespan=All years. Databases=SCI-EXPANDED, SSCI, A&HCI, CPCI-S, CPCI-SSH, BKCI-S, BKCI-SSH, CCR-EXPANDED, IC.
LILACS (BIREME)	Florbetaben OR NEURACEQ OR neuraceq* OR "florbetaben-fluorine-18" OR "18F-BAY94-9172" OR "[18F]Florbetaben" OR "florbetaben-PET" [Words]
CINAHL (EBSCOhost) (1980 to May 2017)	S1 TX Florbetaben S2 TX NEURACEQ S3 TX neuraceq* S4 TX "florbetaben-fluorine-18" S5 TX "18F-BAY94-9172" S6 TX "[18F]Florbetaben" S7 TX "florbetaben-PET" S8 S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7
ClinicalTrials.gov (www.clinicaltrials.gov)	Florbetaben OR NEURACEQ OR neuraceq* OR "florbetaben-fluorine-18" OR "18F-BAY94-9172" OR "[18F]Florbetaben" OR "florbetaben-PET"
World Health Organization International Clinical Trials Registry Platform (WHO ICTRP) (http://apps.who.int/trialsearch)	Florbetaben OR NEURACEQ OR neuraceq OR "florbetaben-fluorine-18" OR "18F-BAY94-9172" OR "[18F]Florbetaben" OR "florbetaben-PET"
ALOIS, the Cochrane Dementia & Cognitive Improvement Group's specialized register of dementia studies (http://www.medicine.ox.ac.uk/alois/)	Imaging AND PET

Appendix 3. Tables (2 × 2) cross-relating index test results of the reference standards

Table 1. Progression from mild cognitive impairment (MCI) to Alzheimer's disease dementia (ADD)

Index test information	References standard information	
	ADD present	ADD absent
Index test-positive	^{18}F -florbetaben PET ligand for $\text{A}\beta$ (+) who progress to ADD (TP)	^{18}F -florbetaben PET ligand for $\text{A}\beta$ (+) who remain MCI (FP) and ^{18}F -florbetaben PET ligand $\text{A}\beta$ (+) who progress to non-ADD (FP)
Index test-negative	^{18}F -florbetaben PET ligand for $\text{A}\beta$ (-) who progress to ADD (FN)	^{18}F -florbetaben PET ligand for $\text{A}\beta$ (-) who remain MCI (TN) and ^{18}F -florbetaben PET ligand for $\text{A}\beta$ (-) who progress to non-ADD (TN)

ADD: Alzheimer's disease dementia

FN: false negative

FP: false positive

MCI: mild cognitive impairment

PET: positron emission tomography

TN: true negative

TP: true positive

Table 2. Progression from mild cognitive impairment (MCI) to non-Alzheimer's disease dementia (non-ADD)

Index test information	References standard information	
	Non-ADD present	Non-ADD absent
Index test-positive	^{18}F -florbetaben PET ligand for $\text{A}\beta$ (+) who progress to non-ADD (TP)	^{18}F -florbetaben PET ligand for $\text{A}\beta$ (+) who remain MCI (FP) and ^{18}F -florbetaben PET ligand for $\text{A}\beta$ (+) who progress to ADD (FP)
Index test-negative	^{18}F -florbetaben PET ligand for $\text{A}\beta$ (-) who progress to non-ADD (FN)	^{18}F -florbetaben PET ligand for $\text{A}\beta$ (-) who remain MCI (TN) and ^{18}F -florbetaben PET ligand for $\text{A}\beta$ (-) who progress to ADD (TN)

ADD: Alzheimer's disease dementia

FN: false negative

FP: false positive

MCI: mild cognitive impairment

PET: positron emission tomography

TN: true negative

TP: true positive

Table 3. Progression from mild cognitive impairment (MCI) to any form of dementia

Index test information	References standard information	
	Any forms of dementia present	Dementia absent

(Continued)

Index test-positive	¹⁸ F-florbetaben PET ligand for Aβ (+) who progress to any form of dementia (TP)	¹⁸ F-florbetaben PET ligand for Aβ (+) who remain MCI (FP)
Index test-negative	¹⁸ F-florbetaben PET ligand for Aβ (-) who progress to any form of dementia (FN)	¹⁸ F-florbetaben PET ligand for Aβ (-) who remain MCI (TN)

FN: false negative

FP: false positive

MCI: mild cognitive impairment

PET: positron emission tomography

TN: true negative

TP: true positive

Appendix 4. Assessment of methodological quality table: Quality Assessment of Diagnostic Accuracy Studies 2 (QUADAS-2) tool

Domain	Patient selection	Index test	Reference standard	Flow and timing
Description	Describe methods of patient selection: describe included patients (prior testing, presentation, intended use of index test and setting)	Describe the index test and how it was conducted and interpreted	Describe the reference standard and how it was conducted and interpreted	Describe any patients who did not receive the index test(s) or reference standard, or both, or who were excluded from the 2 × 2 table (refer to flow diagram): describe the time interval and any interventions between index test(s) and reference standard
Signalling questions (yes/no/unclear)	Was a consecutive or random sample of patients enrolled?	Were the index test results interpreted without knowledge of the results of the reference standard?	Is the reference standard likely to correctly classify the target condition?	Was there an appropriate interval between index test(s) and reference standard?
	Was a case-control design avoided?	If a threshold was used, was it prespecified?	Were the reference standard results interpreted without knowledge of the results of the index test?	Did all patients receive a reference standard?
	Did the study avoid inappropriate exclusions?			Did all patients receive the same reference standard? Were all patients included in the analysis?
Risk of bias (high/low/unclear)	Could the selection of patients have introduced bias?	Could the conduct or interpretation of the index test have introduced bias?	Could the reference standard, its conduct, or its interpretation have introduced bias?	Could the patient flow have introduced bias?
Concerns regarding	Are there concerns that the included pa-	Are there concerns that the index test,	Are there concerns that the target condition as	

(Continued)

applicability (high/low/un- clear)	tients do not match the review question?	its conduct, or inter- pretation differ from the review question?	defined by the reference standard does not match the review question?
--	---	---	---

Appendix 5. Anchoring statements for quality assessment of ^{18}F -florbetaben PET ligand for A β diagnostic studies

Table 4. Review question and inclusion criteria

Category	Review question	Inclusion criteria
Patients	Participants with mild cognitive impairment (MCI), no dementia	Participants that fulfil the criteria for the clinical diagnosis of MCI at baseline
Index test	^{18}F -florbetaben PET ligand for A β bio- marker	^{18}F -florbetaben PET ligand for A β biomarker
Target condition	Alzheimer's disease dementia (ADD) (pro- gression from MCI to ADD) Any other forms of dementia (progres- sion from MCI to any other forms of de- mentia	ADD (progression from MCI to ADD) Any other forms of dementia (progression from MCI to any other forms of dementia)
Reference standard	NINCDS-ADRDA; DSM; ICD; McKeith cri- teria; Lund criteria; International Behav- ioural Variant FTD Criteria Consortium; NINDS-ARIEN criteria	NINCDS-ADRDA; DSM; ICD; McKeith criteria; Lund criteria; International Behavioural Variant FTD Criteria Consortium; NINDS-ARIEN criteria
Outcome	N/A	Data to construct a 2×2 table
Study design	N/A	Longitudinal cohort studies and nested case-control studies if they incorporate a delayed verification design (case-con- trol nested in cohort studies)

ADD: Alzheimer's disease dementia

DSM: Diagnostic and Statistical Manual of Mental Disorders

FTD: Frontotemporal dementia

ICD: International Classification of Diseases

MCI: Mild cognitive impairment

NINCDS-ADRDA: National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association

NINDS-AIREN: National Institute of Neurological Disorders and Stroke and Association Internationale pour la Recherche et l'Enseignement en Neurosciences

PET: Positron emission tomography

Anchoring statements for quality assessment ^{18}F -florbetaben PET ligand for A β diagnostic studies

We have provided some core anchoring statements for quality assessment in the diagnostic test accuracy (DTA) review of the ^{18}F -florbetaben PET ligand for A β biomarker in dementia. These statements are designed for use with the Quality Assessment of Diagnostic Accuracy Studies 2 (QUADAS-2) tool and are based on the guidance for quality assessment of DTA reviews of Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) in dementia (Quinn 2014). In assessing individual items, the score of unclear should only be given if there is genuine uncertainty. In these situations, we contacted the relevant study teams for additional information. Whenever we scored one question as high risk of bias, we considered the study as having a high risk of bias.

Table 5. Anchoring statements to assist with the 'Risk of bias' assessment

Question	Response and weight- ing	Explanation
Patient selection		
Was the sampling method appropriate?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	Where sampling is used, the designs least likely to cause bias are consecutive sampling or random sampling. Sampling that is based on volunteers or selecting subjects from a clinic or research resource is prone to bias.
Was a case-control or similar design avoided?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	Designs similar to case-control that may introduce bias are those designs where the study team deliberately increase or decrease the proportion of subjects with the target condition, which may not be representative. Some case-control methods may already be excluded if they mix subjects from various settings.
Are exclusion criteria described and appropriate?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	<p>We automatically graded the study as unclear if the study authors did not detail exclusions (pending contact with study authors).</p> <p>Where a study details exclusions, we graded the study as 'low risk' if we considered exclusions to be appropriate. Certain exclusions common to many studies of dementia are: medical instability; terminal disease; alcohol/substance misuse; concomitant psychiatric diagnosis; other neurodegenerative conditions.</p> <p>Exclusions are not appropriate if they comprise 'difficult to diagnose' patients.</p> <p>We labelled post-hoc and inappropriate exclusions as at 'high risk' of bias.</p>
Index test		
Was the ¹⁸ F-florbetaben PET ligand for Aβ bio-marker's assessment/interpretation performed without knowledge of clinical dementia diagnosis?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	<p>Terms such as 'blinded' or 'independently and without knowledge of' are sufficient and full details of the blinding procedure are not required. Interpretation of the results of the index test may be influenced by knowledge of the results of the reference standard. If the index test is always interpreted prior to the reference standard, then the person interpreting the index test cannot be aware of the results of the reference standard and so this item could be rated as 'yes'.</p> <p>For certain index tests, the result is objective and knowledge of the reference standard should not influence the result, e.g. level of protein in cerebrospinal fluid; in this instance, the quality assessment may be 'low risk' even if blinding was not achieved.</p>
Was the ¹⁸ F-florbetaben PET ligand for Aβ bio-marker's threshold prespecified?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	For scales and biomarkers, there is often a reference point (in units or categories) above which subjects are classified as 'test-positive'; this may be referred to as the threshold, clinical cut-off, or dichotomisation point. A study is classified at high risk of bias if the study authors define the optimal cut-off post-hoc based on their own study data because selecting the threshold to maximise sensitivity and/or specificity may lead to overoptimistic measures of test performance. Certain papers may use an alternative methodology for analysis that does not use thresholds and these papers should be classified as not applicable.
Was the ¹⁸ F-florbetaben PET ligand for Aβ scan interpretation done by a trained reader physician?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	<p>If a trained reader physician performed the scan interpretation, we scored this item as 'yes'.</p> <p>If no definition of trained reader was done, we scored this item as 'unclear'.</p> <p>If a nontrained reader physician performed the scan interpretation, we scored this item as 'no'.</p>

(Continued)

Did the study provide a clear definition of what was considered to be a ^{18}F -florbetaben PET ligand for A β biomarker's positive result?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	If the study clearly stated the definition of a positive result (e.g. SUV), we scored this item as 'yes'. If the study did not give a definition of what it considered a positive result or the definition of a positive result varied between the participants, we scored this item as 'no'. If the study gave insufficient information to permit judgement, we scored the item as 'unclear'.
Reference standard		
Is the assessment used for clinical diagnosis of dementia acceptable?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	Commonly used international criteria to assist with clinical diagnosis of dementia included those detailed in DSM-IV and ICD-10. Criteria specific to dementia subtypes included but were not limited to NINCDS-ADRDA criteria for Alzheimer's dementia; McKeith criteria for Lewy body dementia; Lund criteria and International Behavioural Variant FTD Criteria Consortium for frontotemporal dementia; and the NINDS-AIREN criteria for vascular dementia. Where the criteria used for assessment were not familiar to the review authors or the Cochrane Dementia and Cognitive Improvement group ('unclear'), we classified this item as 'high risk of bias'.
Were clinical assessments for dementia performed without knowledge of the ^{18}F -florbetaben PET ligand for A β biomarker?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	Terms such as 'blinded' or 'independently and without knowledge of' were sufficient and full details of the blinding procedure were not required. Interpretation of the results of the reference standard may be influenced by knowledge of the results of the index test.
Patient flow		
Was there an appropriate interval between ^{18}F -florbetaben PET ligand for A β biomarker and clinical dementia assessment?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	As we test the accuracy of the ^{18}F -florbetaben PET ligand for A β biomarker for MCI progression to dementia, there will always be a delay between the index test and the reference standard assessments. The time between the reference standard and the index test will influence the accuracy (Geslani 2005; Okello 2007; Visser 2006), and therefore we noted time as a separate variable (both within and between studies) and will test its influence on the diagnostic accuracy. We have set a minimum mean time to follow-up assessment of 1 year. If more than 16% of subjects have assessment for MCI progression before nine months, this item was scored 'no'.
Did all subjects get the same assessment for dementia regardless ^{18}F -florbetaben PET ligand for A β biomarker?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	There may be scenarios where participants who score 'test-positive' on the index test have a more detailed assessment. Where dementia assessment differs between participants, this should be classified as high risk of bias.
Were all patients who received ^{18}F -florbetaben PET ligand for A β biomarker's assessment included in the final analysis?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	If the number of patients enrolled differs from the number of patients included in the 2×2 table, then there is the potential for bias. If patients lost to dropouts differ systematically from those who remain, then estimates of test performance may differ. If there are dropouts, these should be accounted for; a maximum proportion of dropouts for a study to remain at low risk of bias has been specified as 20%.
Were missing ^{18}F -florbetaben PET ligand for	No = high risk of bias Yes = low risk of bias	Where missing or uninterpretable results are reported, and if there is substantial attrition (we have set an arbitrary value of 50% missing data), we will score

(Continued)

Aβ biomarker's results reported?	Unclear = unclear risk of bias	this as 'no'. If the study did not report these results, we scored this as 'unclear' and we contacted the study authors.
Was the study with ¹⁸ F-florbetaben PET ligand for Aβ biomarker free of commercial funding?	No = high risk of bias Yes = low risk of bias Unclear = unclear risk of bias	If the funding source is clearly stated and is not commercial, this should be scored as 'no'. If the funding source is clearly stated and is commercial, this should be scored as 'yes'. If not enough information is given to assess whether the funding source is commercial, the scored is 'unclear'.

Anchoring statements to assist with assessment for applicability

Question	Explanation
Were included patients representative of the general population of interest?	The included patients should match the intended population as described in the review question. The review authors should consider population in terms of symptoms; pretesting; potential disease prevalence; setting. If there is a clear ground for suspecting an unrepresentative spectrum, the item should be rated poor applicability.

Index test

Were sufficient data on ¹⁸ F-florbetaben PET ligand for Aβ biomarker's application given for the test to be repeated in an independent study?	Variation in technology, test execution, and test interpretation may affect estimate of accuracy. In addition, the background, and training/expertise of the assessor should be reported and taken in consideration. If ¹⁸ F-florbetaben PET ligand for Aβ biomarker was not performed consistently, this item should be rated poor applicability.
--	---

Reference standard

Was clinical diagnosis of dementia made in a manner similar to current clinical practice?	For many reviews, inclusion criteria and 'Risk of bias' assessments will already have assessed the dementia diagnosis. For certain reviews, an applicability statement relating to the reference standard may not be applicable. There is the possibility that a form of dementia assessment, although valid, may diagnose a far larger proportion of people with disease than usual clinical practice. In this instance, the item should be rated poor applicability.
---	--

DSM: Diagnostic and Statistical Manual of Mental Disorders

FTD: Frontotemporal dementia

ICD: International Classification of Diseases

MCI: Mild cognitive impairment

NINCDS-ADRDA: National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association

NINDS-AIREN: National Institute of Neurological Disorders and Stroke and Association Internationale pour la Recherche et l'Enseignement en Neurosciences

PET: Positron emission tomography

CONTRIBUTIONS OF AUTHORS

Gabriel Martínez, Robin WM Vernooij, and Paulina Fuentes Padilla: contributed to conception, design, and draft of the protocol; overall responsibility of study selection; data extraction; contacted the authors; draft of discussion and authors' conclusion sections.

Javier Zamora: reviewed draft protocol, updated statistical methods section, performed statistical analyses and final manuscript.

Leon Flicker: contributed to conception, and designed and reviewed draft protocol and final manuscript.

Xavier Bonfill Cosp: reviewed draft protocol and final manuscript.

DECLARATIONS OF INTEREST

Gabriel Martínez has no known conflicts of interest.
Robin WM Vernooij has no known conflicts of interest.
Paulina Fuentes Padilla has no known conflicts of interest.
Javier Zamora has no known conflicts of interest.

Leon Flicker has no known conflicts of interest.
Xavier Bonfill Cosp has no known conflicts of interest.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- National Institute for Health Research (NIHR), UK.

This protocol was supported by the NIHR, via Cochrane Infrastructure funding to the Cochrane Dementia and Cognitive Improvement group. The views and opinions expressed therein are those of the protocol authors and do not necessarily reflect those of the Systematic Reviews Programme, the NIHR, the NHS, or the Department of Health.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

- We added as the reference standard the definition of progressive supranuclear palsy (PSP) ([Hauw 1994](#)).